



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Original Article

Herbivores Coprolites from Chehrabad Salt Mine of Zanjan, Iran (Sassanid Era, 224-651 AD) Reveals Eggs of Strongylidae and Anoplocephalidae Helminths

Masoumeh MEIGOUNI ¹, *Mahsasadat MAKKI ², Ali HANILOO ¹, Zeynab ASKARI ², Iraj MOBEDI ², Saied Reza NADDAF ³, Nicole BOENKE ⁴, Thomas STOLLNER ⁴, Abolfazl AALI ⁵, Zahra HEIDARI ⁶, *Gholamreza MOWLAVI ^{2,7}

1. Department of Parasitology and Mycology, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran
2. Department of Medical Parasitology and Mycology, School of Public Health, Tebran University of Medical Sciences, Tebran, Iran
3. Department of Parasitology, Pasteur Institute of Iran, Tebran, Iran
4. Ruhr Universität Bochum, Institut für Archäologische Wissenschaften, Am Bergbaumuseum 31, D-44791 Bochum, Germany
5. Archaeological Museum of Zanjan, Zanjan, Iran
6. Department of Medical Microbiology and Parasitology, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran
7. Center for Research of Endemic Parasites of Iran (CREPI), Tebran University of Medical Sciences, Tebran, Iran

Received 14 Mar 2019

Accepted 26 Jun 2019

Keywords:

Paleoparasitology;
Herbivores;
Strongyle;
Anoplocephalan;
Iran

***Correspondence**

Emails:

Molavig@yahoo.com
makki62@gmail.com

Abstract

Background: The ancient Chehrabad Salt mine, a well-known archaeological site in Iran, has recently received increasing interest from Iranian and international archeologists. Also, the biological remains from this site have provided valuable sources for studying the pathogenic agents of ancient times. This study aimed to identify the parasitic helminth eggs preserved in the herbivores coprolites.

Methods: From 2011 to 2015, we received three coprolites belonging to herbivorous animals recovered during excavations in Chehrabad Salt mine of Zanjan, Iran. The coprolites were dated back to the Sassanid era (224-651 AD) by using radiocarbon accelerator mass spectrometry (AMS) and archeological stratigraphy methods. Following rehydration of the specimens in a 0.5% trisodium phosphate solution, the suspensions were mounted in glycerin jelly on glass slides and examined by a light microscope with 100x and 400x magnifications.

Results: Two coprolites belonged to donkeys and one to an unknown herbivore species. The recovered eggs belonged to members of two helminths families, Strongylidae, and Anoplocephalidae. Also, within the two coprolites, some mites, presumably of the order Oribatida, were observed.

Conclusion: The presence of two different nematodes in the equids coprolites provide clues of the burden of helminths infection on working animal at the Sassanid time and demonstrates the appropriate preservation condition of biological remains in the ancient salt mine of Chehrabad as well.

Introduction

The parasites in the ancient biological remains provide an image of the status of parasitic infections in a specific period in the past. The coprolites are a valuable source for tracing parasitic agents in humans and animals over time. They can as well shed light on the emergence and elimination of parasitic infections in time and provide clues on the intercontinental migrations of humans and animals (1).

The parasitic worms of equids such as strongyles have a worldwide distribution (2-6), while the available data of their presence in ancient times is very scarce (7, 8). Strongyle nematodes are amongst the most prevalent pathogenic helminth inhabiting the large intestine of herbivores. These nematodes are soil-transmitted helminths and are acquired by the animals via the ingestion of the third-stage larvae (L3). Infection with this parasite produces mild to severe clinical symptoms due to the migration of the larvae (9, 10). In contrast to strongyles, the Anoplocephalidae cestodes are arthropod-borne helminths and infect the hosts via the ingestion of oribatid mites (11). Previously, in Chehrabad salt mine archeological site, eggs of several helminths that infect humans, rodents, and carnivores were identified (12-14). The appropriate preservation condition in this ancient site has preserved biological remains over the past millennia. The present study describes the identification of the helminth eggs in herbivores coprolites recovered from Chehrabad Salt mine archeological site.

Materials and Methods

Samples

From 2011 to 2015, we received three coprolites (Code numbers: 2605-286, 2357-282, and 2462-124) from the Cultural Heritage Organization of Zanjan, northwestern Iran. Of three coprolites, two (code numbers 2605-286

and 2462-124) matched donkey droppings based on the size and their typical round (cubic) kidney shape. The third sample that had lost its original shape was attributed to a herbivore by archaeobotanical analysis.

The coprolites were dated back to the Sassanid era (224-651AD) by using radiocarbon accelerator mass spectrometry (AMS) and archeological stratigraphy methods as described elsewhere (15,16). The samples were kept in the Laboratory of Helminthology at the School of Public Health, Tehran University of Medical Sciences, until used.

Microscopical examination

The samples were rehydrated in a 0.5% trisodium phosphate solution (Na_3PO_4) for one week (17, 18). The suspension from each specimen was mounted in glycerin jelly on 200 glass slides and examined under a light microscope at 100X and 400X magnifications. The retrieved eggs were photographed by a camera-equipped microscope (Labomed LX 500, Springfield, New Jersey, USA); their measurements were recorded and compared with similar eggs available in the literature. The helminths' eggs were identified based on the measurements and morphological characters available in taxonomic keys (19, 20).

Results

Based on reliable morphological features and measurements, we identified two types of eggs belonging to the members of the families Strongylidae and Anoplocephalidae (Fig.1). The typical thick shells with the pyriform apparatus inside, demonstrated Anoplocephalidae eggs (Fig1. A&B), and the thin oval shell, containing larvae, clearly represented the strongyles' eggs (Fig.1, C&D) (19, 20).

The specimen with the code number 2605-286 contained both types of helminths eggs, while the specimens with the code numbers

2357-282 and 2462-124 harbored only the eggs of Strongylidae and Anoplocephalidae, respectively (Table 1). Also, within the two coprolites with code numbers 2605-286 and 2462-124, mites of the class Arachnida, subclass Acari were observed. (Fig.2).




Fig. 1: Recovered helminth eggs from donkeys (A, B&C) and unidentified herbivore (D) coprolites. Strongylidae eggs (C&D) with larvae inside, and Anoplocephalidae eggs (A&B).



Fig. 2: The mite in herbivores coprolites recovered in Chehrabad Archeological Site, Zanjan, Iran.

Table 1: Helminth eggs retrieved from the herbivores coprolites

<i>Code number</i>	<i>Anoplocephalidae eggs</i>	<i>Strongylidae eggs</i>
2605-286	+	+
2357- 282	-	+
2462-124	+	-
Egg measurements (μm)	53.6 ± 3.2	72.30 ± 7.09
(length and width range mean)	53.4 ± 1.9	37.35 ± 5.53

Discussion

The coprolites recovered from archaeological sites are valuable sources for identification of the parasites that were prevalent in ancient times; however, well-preserved coprolites are not much available (7, 8). The present study reports the identification of two helminths eggs in the well-preserved ungulate coprolites from Chehrabad salt mine of Zanjan, North

West of Iran. The eggs belonged to the members of families, Strongylidae and Anoplocephalidae, today commonly found in the digestive tracts of the equines.

The presence of mites within the coprolites draws the attention to oribatid mites that might serve as intermediate hosts of *Anoplocephala* tapeworms of cattle or other ruminants. These mites belong to the superorder Acariformes and commonly occur in soil and hu-

mus. Today, equine strongylosis is highly prevalent worldwide (3, 21, and 22) and poses a serious threat to the populations of horses and donkeys (4, 10). In Iran, in Marand, an area adjacent to the Chehrabad Archeological Site, the prevalence rate of strongyle nematodes among 58 donkeys was 100% (23, 24). The occurrence of strongyles among the equids does not pose a public health threat; however, very few reports on human infections with *Bertiella studeri*, a member of the Anoplocephalidae family, are available (25-27).

The infection with some strongyle species, e.g., *S. vulgaris*, may result in the mesenteric arterial obstruction in donkeys (5, 28). Hence, equine strongylosis in the Sassanid era might have been a significant veterinary problem or even a fatal disease of working animals; the only means of transportation in ancient times (29). Individual animals commonly harbor multiple species, which may apply to the coprolites we examined; precise identification of the species by morphological features of eggs and early-stage larvae is impossible. The interspecific differences in rDNA sequences of *Strongylus* spp. allows the development of reliable molecular tools to differentiate individual eggs (30-33). In addition, the capability of the real-time PCR assay, which detects equivalents of 0.5 strongyle eggs with no cross-reactivity (34), increases the chance of finding helminth eggs in a limited amount of coprolites.

Molecular characterization of the ancient DNA of helminths provides a comparison of the archaic and modern populations of the helminth parasites and elucidates the evolution of parasites over millennia and the coevolution of host and parasite (35, 36).

Conclusion

The presence of two different helminth eggs in the coprolites of equids provide clues of the burden of helminths infection on working animal at the Sassanid time and demonstrates the appropriate preservation condition of biological

remains in the ancient salt mine of Chehrabad.

Acknowledgements

This study was funded by the Center for Research of Endemic Parasites of Iran (CREPI), Tehran University of Medical Sciences (grant No. 17195-160-01-91). We thank our colleagues in the Bochum University, Germany, and the Cultural Heritage, Handicraft and Tourism Organization of Zanjan, Iran, for providing the samples and their scientific collaboration. The authors also acknowledge Ms. Niloofar Paknezhad for her technical assistance.

Conflict of interest

The authors declare there are no issues with this article to be perceived as a conflict of interest.

References

1. Sianto L, Chame M, Silva CS et al. Animal helminths in human archaeological remains: a review of zoonoses in the past. *Rev Inst Med Trop Sao Paulo*. 2009;51(3):119-30.
2. Ayele G, Fescha G, Bojia E, Joe A. Prevalence of gastro-intestinal parasites of donkeys in Dugda Bora District, Ethiopia. *Livestock Research for Rural Development*. 2006; 18(10):14-21.
3. Bu Y, Niu H, Gasser R et al. Strongyloid nematodes in the caeca of donkeys in Henan Province, China. *Acta Parasitologica*. 2009;54(3):263-8.
4. Umur Ş, Acici M. A survey on helminth infections of equines in the Central Black Sea region, Turkey. *Turk J Vet Anim Sci*. 2009;33(5):373-8.
5. Morgan SJ, Stromberg PC, Storts RW et al. Histology and morphometry of *Strongylus vulgaris*-mediated equine mesenteric arteritis. *J Comp Pathol*. 1991;104(1):89-99.
6. Whitlock J, Leasure E. Studies upon *Strongylus vulgaris*. I. The incidence of *Strongylus vulgaris* in

- mid-continental North America and the reaction of the infested ceca. *Am J Hyg.* 1939;29(3):83-7.
7. Le Bailly M, Lepetz S, Samashev Z et al. Palaeoparasitological study of gastro-intestinal content in horses at a Scythian kurgan (3rd century BC) found in Kazakhstan. *Anthropozoologica.* 2008;43(2):69-75.
 8. Dufour B, Hugot JP, Lepetz S, Le Bailly M. The horse pinworm (*Oxyuris equi*) in archaeology during the Holocene: Review of past records and new data. *Infect Genet Evol.* 2015;33:77-83.
 9. Khan M, Roohi N, Rana M. Strongylosis in equines: a review. *J Anim Plant Sci.* 2015;25:1-9.
 10. Shite A, Admassu B, Abere A. Large strongyle Parasites in Equine: A Review. *Adv Biol Res.* 2015;9(4):247-52.
 11. Denegri G, Bernadina W, Perez-Serrano J et al. Anoplocephalid cestodes of veterinary and medical significance: a review. *Folia Parasitol (Praha).* 1998;45(1):1-8.
 12. Nezamabadi M, Mashkour M, Aali A et al. Identification of *Taenia* sp. in a natural human mummy (third century BC) from the Chehrabad salt mine in Iran. *J Parasitol.* 2013;99(3):570-2.
 13. Mowlavi G, Makki M, Mobedi I et al. Paleoparasitological findings from rodent coprolites dated at 500 CE Sassanid Era in archeological site of Chehrabad (Douzlakh), salt mine Northwestern Iran. *Iran J Parasitol.* 2014;9(2):188-93.
 14. Mowlavi G, Makki M, Heidari Z et al. *Macracanthorhynchus hirudinaceus* eggs in canine coprolite from the Sasanian Era in Iran (4th/5th Century CE). *IJP.* 2015;10(2):245.
 15. Aali A, Stöllner T. (Eds.). The archaeology of the salt miners. *Interdisciplinary Research 2010–2014.* Deutsches Bergbau-Museum;2015.
 16. Aali A, Abar A, Boenke N, Pollard M, Rühli F, Stöllner T. Ancient salt mining and salt men: the interdisciplinary Chehrabad Douzlakh project in North-Western Iran. *Antiquity.* 2012;86(333).
 17. Reinhard KJ, Confalonieri UE, Herrmann B et al. Recovery of parasite remains from coprolites and latrines: aspects of paleoparasitological technique. *Homo.* 1986;37(4):217-239.
 18. Makki M, Dupouy-Camet J, Sajjadi SMS et al. First Paleoparasitological Report on the Animal Feces of Bronze Age Excavated from Shahr-e Sukhteh, Iran. *Korean J Parasitol.* 2017;55(2):197-201.
 19. Soulsby EJJL. Helminths, arthropods and protozoa of domesticated animals: Bailliere Tindall; 1982.
 20. Kaufmann J. Parasitic infections of domestic animals: a diagnostic manual. Birkhäuser Basel. 1996.
 21. Getachew M, Trawford A, Feseha G, Reid S. Gastrointestinal parasites of working donkeys of Ethiopia. *Trop Anim Health Prod.* 2010;42(1):27-33.
 22. Matthee S, Krecek RC, Milne SA. Prevalence and biodiversity of helminth parasites in donkeys from South Africa. *J Parasitol.* 2000;86(4):756-62.
 23. Tavassoli M, Yamchi JA, Hajipour N. A survey on the prevalence of strongyles species in working donkeys in North-West of Iran. *J Parasit Dis.* 2016;40(4):1210-2.
 24. Eslami A, Kiai B. Identification of cyathostomes in equines in Iran. *Iran J Vet Res* 2007;8(1):45-57.
 25. Bhagwant S. Human *Bertiella studeri* (family Anoplocephalidae) infection of probable Southeast Asian origin in Mauritian children and an adult. *Am J Trop Med Hyg.* 2004;70(2):225-8.
 26. Adams A, Webb L. Two further cases of human infestation with *Bertiella studeri* (Blanchard, 1891) Stiles and Hassall, 1902, with some observations on the probable synonymy of the specimens previously recorded from man. *Ann Trop Med Parasitol.* 1933;27(3):471-5.
 27. Sun X, Fang Q, Chen XZ et al. *Bertiella studeri* infection, China. *Emerg Infect Dis.* 2006;12(1):176
 28. Borji H, Moosavi Z, Ahmadi F. Cranial Mesenteric Arterial Obstruction Due To *Strongylus vulgaris* Larvae in a Donkey (*Equus asinus*). *Iran J Parasitol.* 2014;9(3):441-4.
 29. Bendrey R. Animal paleopathology. *Encyclopedia of Global Archaeology.* 2014:258-65.
 30. Ampbell AJ, Gasser RB, Chilton NB. Differences in a ribosomal DNA sequence of

- Strongylus* species allows identification of single eggs. Int J Parasitol. 1995;25(3):59–65.
31. Gasser RB, Stevenson LA, Chilton NB, Nansen P, Bucknell DG, Beveridge I. Species markers for equine strongyles detected in intergenic rDNA by PCR-RFLP. Mol Cell Probes. 1996;10(5):371–8.
 32. Hung GC, Gasser RB, Beveridge I, Chilton NB. Species-specific amplification by PCR of ribosomal DNA from some equine strongyles. Parasitology. 1999;119:69–80.
 33. Zhang LP, Hu M, Chilton NB, Huby-Chilton F, Beveridge I, Gasser RB. Nucleotide alterations in the D3 domain of the large subunit of ribosomal DNA among 21 species of equine strongyle. Mol Cell Probes. 2007;21(2):111–5.
 34. Andersen UV, Howe DK, Olsen SN, Nielsen MK. Recent advances in diagnosing pathogenic equine gastrointestinal helminths: the challenge of prepatent detection. Vet Parasitol. 2013;192(1-3):1-9.
 35. Leles D, Araújo A, Ferreira LF, Vicente ACP, Iñiguez AM. Molecular paleoparasitological diagnosis of *Ascaris* sp. from coprolites: new scenery of ascariasis in pre-Columbian South America times. Mem Inst Oswaldo Cruz. 2008;103(1):106-108.
 36. Iñiguez AM, Reinhard KJ, Araújo A, Ferreira LF, Vicente ACP. *Enterobius vermicularis*: ancient DNA from North and South American human coprolites. Mem Inst Oswaldo Cruz. 2003;98:67-69.