



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Original Article

Molecular Detection of *Dirofilaria immitis* Specific Gene from Infected Dog Blood Sample Using Polymerase Chain Reaction

In Young OH¹, Kyung Tae KIM², *Ho Joong SUNG^{1,3}

1. Dept. of Biomedical Laboratory Science, College of Health Sciences, Eulji University, Seongnam-si, Gyeonggi-do, Korea
2. ALPHAGENE Co., Ltd., Singu University, Business Incubation Center, Seongnam, Korea
3. Dept. of Senior Healthcare, BK21 Plus Program, Graduate School, Eulji University, Daejeon, Korea

Received 21 Sep 2016
Accepted 18 Mar 2017

Keywords:
Dirofilaria immitis,
COI,
Molecular diagnosis,
PCR

***Correspondence
Email:**
hjsung@eulji.ac.kr

Abstract

Background: *Dirofilaria immitis*, a filarial nematode, is the most important parasite-affecting dogs, causing cardiopulmonary dirofilariasis. Current diagnostic tools for detecting *D. immitis* include morphological assays, antigen detection, and X-ray. Herein, we developed a method for the molecular detection of *D. immitis* in blood using polymerase chain reaction (PCR).

Methods: The study was conducted at Eulji University, Republic of Korea in 2016. To detect *D. immitis*-specific gene regions, we aligned the cytochrome c oxidase subunit I (COI) genes of seven filarial nematodes and designed primers targeting the unique region. We used dog glyceraldehyde-3-phosphate dehydrogenase (GAPDH)-targeted primers as the internal control. We conducted PCR-amplified genomic DNA from canine blood samples. The products were confirmed by sequencing.

Results: Gene alignment revealed a *D. immitis* COI-specific gene region, and the activity of designed primers was confirmed by PCR and sequencing. Plasmid DNA made from the PCR products was a positive control. The limit of detection for our method was 50 copies. The *D. immitis* COI and dog GAPDH genes could be discriminated from blood samples simultaneously.

Conclusion: This study provides a method for highly specific and sensitive molecular diagnosis of *D. immitis* used as a diagnostic and therapeutic tool from the early stage of infection.

Introduction

Dirofilaria immitis is a filarial nematode that causes heartworm disease in canines, felines, various wild mammals, and some human populations with in-

creasing incidence in tropical, subtropical, and some temperate regions (1-6). Mosquitoes are the most important vectors for accidental infection of *D. immitis*. When infected mosquito

bites a susceptible animal, the third-stage (L3) larvae of *D. immitis* in the head of the mosquito are transmitted to the new final host. After infection, the larvae grow in the blood to the adult stage and subsequently are transferred to the heart of the host, where they cause pulmonary dirofilariasis (1, 2, 4, 7, 8).

Current methods for diagnosis of *D. immitis* include microscopy-based morphological assays, immunochromatographic antigen detection or ELISA, and X-ray imaging of the main pulmonary artery and the right side of the heart. Methods for molecular detection by PCR are also under development (7, 9-11). Among these diagnostic tools, the most widely used method is microscopic morphological examination of microfilariae from blood samples. However, the number of circulating microfilariae does not correlate with the number of adult heartworms and therefore does not indicate disease severity. Furthermore, this approach has limited sensitivity, and expert analysis is required to distinguish among filarial parasite species because of their rather similar morphology (7, 12). In addition, antigen detection targets the antigens released from the reproductive tract of adult female worms and can give false-negative results during the first 5–8 months of infection due to low worm counts, immature infections, and all-male infections (8,10). Therefore, the development of molecular diagnostic techniques for early diagnosis and therapeutic monitoring is essential, and identification of a sensitive diagnostic molecular marker for heartworm infections is crucial for controlling the disease (7, 13).

This study aimed to develop a PCR-based molecular detection method targeting a *D. immitis*-specific gene in the peripheral blood of infected dogs.

Materials and Methods

Identification of the specific gene region of *D. immitis*

The study was conducted at Eulji University, Republic of Korea in 2016. To search for the

specific gene region of *D. immitis*, the seven filarial cytochrome c oxidase subunit I (COI) gene sequences available were multiple aligned using the Align Sequences Nucleotide BLAST (Basic local alignment search tool) program (NCBI);

https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&PAGE_TYPE=BlastSearch&LINK_LOC=blasthome. At the web page, enter the accession number of *D. immitis* COI (EU159111.1) in the Query Sequence box, and then enter the accession number of *Setaria tundra* COI (AJ544874.1), *Setaria digitate* COI (EF174428.1), *Brugia malayi* COI (AJ271610.1), *Wuchereria bancrofti* COI (AJ271612.1), and *Onchocerca volvulus* COI (AM749284.1) in the Subject Sequence box.

The multiple alignments can be obtained by clicking the [BLAST] button at the bottom of the web page. The genome sequences of COI genes of filarial nematodes were obtained from GenBank. Compared with *D. immitis* COI, the six other filarial COI gene sequences showed nucleotide identities as follows; *D. repens* COI, 90%; *S. tundra* COI, 89%; *S. digitate* COI, 88%; *B. malayi* COI, 85%; *W. bancrofti* COI, 86%; and *O. volvulus* COI, 89%. The BLAST tree of these results is presented in Fig. 1.

The BLAST tree was created by clicking the [distance tree of result] button at the bottom of the result page of multiple alignments and using the Radial Tree layout.

The BLAST phylogenetic tree for the COI gene was generated using in silico coverage analysis on the NCBI website. The identities (%) of these species compared with *D. immitis* are reported in the text.

Design of primers

The genome sequences of *D. immitis* and *Canis lupus familiaris* (dog) were obtained from GenBank. Based on the multiple sequence alignment described above, primers were designed to target the *D. immitis*-specific gene region (Fig. 2).




Fig. 1: BLAST tree of filarial nematode COI genes

Primers targeting specific gene regions were designed based on a highly conserved region of the COI gene of *D. immitis* to amplify a 150-bp fragment (forward: ATT GGG TGC CCC TGA AAT GG; reverse: CCC TCT ACA CTC AAA GGA GGA) and the glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) gene of dog to amplify a 106-bp fragment (forward: CAT GTT TGT GAT GGG CGT GAA; reverse: GAT GAC TTT GGC TAG AGG AGC). The primer specificities were evaluated using BLASTN 2.3.1+ on the NCBI website. All primers used in this study were synthesized by Integrated DNA Technologies (Coralville, IA, USA).

The boxed region in the top line shows the forward primer sequence of the *D. immitis* COI, and that in the second line shows the reverse primer sequence of the gene. The query was *D. immitis* COI (GenBank: EU159111.1), and the subjects are *Dirofilaria repens* COI (AB973225.1), *Setaria tundra* COI (AJ544874.1), *Setaria digitata* COI (EF174428.1), *Brugia malayi* COI (AJ271610.1), *Wuchereria bancrofti* COI

(AJ271612.1), and *Onchocerca volvulus* COI (AM749284.1).

Preparation of genomic DNA from dog blood samples

D. immitis-infected blood samples isolated from random infected source dogs were gifted from Seoul National University, Republic of Korea.

The samples were from Beagles and collected in Jul 2015. Blood samples from healthy volunteers and uninfected dogs that received a routine complete blood count (CBC) test in Veterinary Medical Teaching Hospital of Seoul National University were used as negative controls. Genomic DNA from blood samples collected in EDTA (ethylenediaminetetraacetic acid) tubes was extracted using the QIAamp DNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. The concentration of genomic DNA was determined with a NanoDrop spectrophotometer (ThermoFisher Scientific, Sunnyvale, CA, USA).




Fig. 2: Primer design based on alignment results

PCR amplification

The PCR mixture was prepared with 1× Master Mix (Alphagene, Republic of Korea), 50 ng genomic DNA, and 500 nM primers. The PCR protocol included an initial denaturation step at 95 °C for 5 min; 35 cycles of denaturation at 95 °C for 20 sec, annealing at 60 °C for 20 sec, and extension at 72 °C for 40 sec; and final elongation for 7 min at 72 °C. All amplification reactions were performed on a Veriti™ Dx 96-well Thermo Cycler (Applied Biosystems, Foster City, CA, USA) in a total

volume of 20 µL. PCR products were analyzed by electrophoresis on 2% agarose gels.

DNA sequencing and cloning

To confirm primer activity, the PCR products were sequenced (Bionics, Republic of Korea). After confirmation of the sequences, the PCR products were purified and inserted into pLUG-Prime® TA-cloning vectors (iNtRON Biotechnology, Republic of Korea). The plasmid DNA (pDNA) was cloned and used as a positive control for the PCR reaction. The sequencing results are presented in Table 1.

Table 1: Sequencing results

	Description	Query cover	E value	Ident	Accession
1	<i>Dirofilaria immitis</i> isolate HU11 cytochrome c oxidase subunit I (COI) gene, partial cds; mitochondrial	100%	2E-69	100%	KM452920.1
2	<i>Canis lupus familiaris</i> glyceraldehyde-3-phosphate dehydrogenase (GAPDH), mRNA	100%	1E-45	100%	NM_001003142.2

Dual PCR amplification

To detect *D. immitis* COI and dog *GAPDH* simultaneously, the working concentrations of the primers were optimized. The dual PCR condition was the same as that described above.

Results

Primer activity confirmation

To confirm the activity of primers, dog blood samples were PCR amplified with the primers, and the PCR products were sequenced. The sequences of the PCR products exactly matched those of the *D. immitis* COI and the dog *GAPDH* gene regions (data not shown). To use the amplicons as positive controls, pDNAs were constructed and cloned using TA-cloning vector. The pDNA was the-

reafter sequenced to verify the insertion and amplified using PCR (Fig. 3).

Lanes 1–3 (150 bp) show the insert of the *D. immitis* COI pDNA, and lanes 4–5 (106 bp) show that of dog *GAPDH*. Lane 0: 100-bp DNA ladder.

Limit of detection

To determine the limit of detection, *D. immitis* pDNA was serially diluted from 5×10^7 to 5×10^1 copies and detected by PCR (Fig. 4). The DNA copy number was estimated from the calculated molecular weight of the *D. immitis* pDNA:

$$\text{copy number} = \frac{\text{amount of dsDNA (ng)} \times 3.0221 \times 10^{23} \text{ (molecules/mole)}}{\text{length of dsDNA} \times 660 \text{ (g/mole)} \times 1 \times 10^9 \text{ (ng/g)}}$$

The detection limit was ≤ 50 copies.




Fig. 3: PCR products of plasmid DNA




Fig. 4: Limit of detection of *D. immitis* DNA

Lanes 1–6: *D. immitis* pDNA; lane 7: distilled water. Ten-fold serial dilutions from 5×10^6 to 50 copies were employed for lanes 1–6, respectively. Lane 0: 100-bp DNA ladder.

Optimization of primer working concentrations

To amplify *D. immitis* COI gene and dog GAPDH gene in the same tube simultaneously, the working concentrations of the primers were optimized (Fig. 5).




Fig. 5: Optimization of primer working concentrations

The optimized primer concentration ratio was 2:1 (500 nM *D. immitis* COI primers to 250 nM dog GAPDH primers). The other concentration ratios yielded strong amplification of one gene but weak amplification of the other due to competition between the two primer sets.

Lanes 1–3: products of dual PCR; lane 4: *D. immitis* COI (500 nM primers); lane 5: dog GAPDH (500 nM primers); lane 6: combined products (1:1) from lanes 4 and 5. The primer concentration ratios (*D. immitis* COI to dog GAPDH) were 1:1 (lane 1), 2:1 (lane 2), and 4:1 (lane 3).

Molecular detection of *D. immitis* from dog blood samples

Using the optimized dual PCR conditions, genomic DNA extracted from dog blood

samples was amplified. *D. immitis* COI pDNA was used as positive DNA and dog GAPDH pDNA was used as an internal control DNA of the assay. *D. immitis* COI pDNA (5×10^6 copies) and dog GAPDH pDNA (5×10^5 copies) were used positive control templates; dog GAPDH pDNA (5×10^5 copies) was used as the negative control template for the *D. immitis* detection assay. In PCR with *D. immitis* COI and dog GAPDH primers, the products of *D. immitis* negative control and negative samples showed only the band for dog GAPDH (106 bp), whereas the products of positive control and positive samples showed two bands for *D. immitis* COI (150 bp) and dog GAPDH; the NTC (no template control) had no band (Fig. 6).

Discussion

A direct relationship was reported between *D. immitis* larval development in the mosquito and temperatures in the range of 18–34 °C, indicating that ambient temperature is the major limiting factor in *D. immitis* transmission in most localities (1, 14). The dog blood samples used in this study were collected in the summer in a high-temperature climate (Republic of Korea). In the Republic of Korea, many people breed pet dogs and administer heartworm medication with or without infection to prevent death due to cardiopulmonary dirofilariasis. However, drug use without pre-diagnosis can lead to development of antibiotic resistance (15). Therefore, a method for accurate and accessible analysis is needed. However, current widely used diagnostic tools have many limitations. To overcome these limitations and improve clinical pathogen detection, we developed a molecular diagnostic tool in this study.

To develop this method, we focused on the mitochondrial COI gene. Mitochondrial DNA (mtDNA) genes have high copy numbers, allowing recovery of large amounts of mtDNA from trace samples compared to nuclear DNA.




Fig. 6: Molecular detection of *D. immitis* using dual PCR amplification. Lanes 1–4: *D. immitis*-negative samples (gDNA from healthy dog blood); lane 5: *D. immitis*-negative control (dog *GAPDH* pDNA as an internal control of *D. immitis* detection); lanes 6–9: *D. immitis*-positive samples (gDNA from infected dog blood); lane 10: positive control (*D. immitis* *COI* pDNA and dog *GAPDH* pDNA); lane 11: NTC (no template control)

MtDNA is maternally inherited. It generally does not undergo recombination; thus, its sequence is identical for all maternally linked relatives, and transmission of mtDNA is consistent across many generations. As a mtDNA gene, *COI* is the so-called “barcode” for identifying the last indicator digit of species diversity. “*COI* may be matched by other mitochondrial genes in its efficacy in resolving such cases of recent divergence. This gene is more likely to provide deeper phylogenetic insights than alternatives such as cytochrome b because changes in its amino acid sequence occur more slowly than those in other mitochondrial genes” (16-20).

In this study, we first identified a *D. immitis*-specific region of *COI*. The sequence identity among *COI* genes from filarial species was 85%–90%, with some sequence regions exhibiting complete identity; however, other regions were rarely the same between species. In the inconsistent region, primers were designed to target specifically the *D. immitis* *COI*. The nucleic acid-based amplification assays were highly sensitive with a limit of detection of fewer than 50 copies and are expected to be highly specific to the *D. immitis* *COI*. The assay also included amplification and detection of

the dog *GAPDH* gene as an internal control. Using adjusted PCR conditions for the duplex reaction, *D. immitis* *COI*, and dog *GAPDH* gene could be detected simultaneously. This assay provides an opportunity to discriminate *D. immitis* from infected dog blood samples in the early stage of infection, and our qualitative PCR results are the first step to enabling future development studies.

Conclusion

This study provides valuable information for future studies of *D. immitis* infection, and it could facilitate investigation of the pathogenesis of *D. immitis*. However, in accordance with the characteristics of end-point PCR, this assay can determine only whether a sample is infected. For more precise therapeutic monitoring, therefore, additional development of the quantitative analysis is needed, and this method warrants further investigation.

Acknowledgements

The Small and Medium Business Administration financially supported this work with grant number, C0300247. We thank Prof.

Byeong Chun Lee, Seoul National University, Republic of Korea for providing dog blood samples.

Conflict of Interests

The authors declare that there is no conflict of interest.

References

1. Watts KJ, Courteny CH, Reddy GR. Development of a PCR- and probe-based test for the sensitive and specific detection of the dog heartworm, *Dirofilaria immitis*, in its mosquito intermediate host. *Mol Cell Probes*. 1999;13(6):425-30.
2. Kronefeld M, Kampen H, Sassnau R et al. Molecular detection of *Dirofilaria immitis*, *Dirofilaria repens* and *Setaria tundra* in mosquitoes from Germany. *Parasit Vectors*. 2014;7:30.
3. Khedri J, Radfar MH, Borji H et al. Canine heartworm in Southeastern of Iran with review of disease distribution. *Iran J Parasitol*. 2014;9(4):560-7.
4. Fu Y, Lan J, Wu X et al. Expression of translationally controlled tumor protein (TCTP) gene of *Dirofilaria immitis* guided by transcriptomic screening. *Korean J Parasitol*. 2014; 52(1):21-6.
5. Rossi MI, Aguiar-Alves F, Santos S et al. Detection of Wolbachia DNA in blood from dogs infected with *Dirofilaria immitis*. *Exp Parasitol*. 2010;126(2):270-2.
6. Nuchprayoon S, Junpee A, Poovorawan Y et al. Detection and differentiation of filarial parasites by universal primers and polymerase chain reaction-restriction fragment length polymorphism analysis. *Am J Trop Med Hyg*. 2005; 73(5):895-900.
7. Rojas A, Rojas D, Montenegro VM et al. Detection of *Dirofilaria immitis* and other arthropod-borne filarioids by an HRM real-time qPCR, blood-concentrating techniques and a serological assay in dogs from Costa Rica. *Parasit Vectors*. 2015;8:170.
8. Schnyder M, Deplazes P. Cross-reactions of sera from dogs infected with *Angiostrongylus vasorum* in commercially available *Dirofilaria immitis* test kits. *Parasit Vectors*. 2012;5:258.
9. Xu D, Zhang J, Shi Z et al. Molecular detection of vector-borne agents in dogs from ten provinces of China. *Parasit Vectors*. 2015;8:501.
10. Oi M, Sato Y, Nakagaki K et al. Detection of *Dirofilaria immitis* DNA in host serum by nested PCR. *Parasitol Res*. 2015;114(10):3645-8.
11. Cardoso L, Mendão C, Madeira de Carvalho L. Prevalence of *Dirofilaria immitis*, *Ehrlichia canis*, *Borrelia burgdorferi* sensu lato, *Anaplasma* spp. and *Leishmania infantum* in apparently healthy and CVBD-suspect dogs in Portugal—a national serological study. *Parasit Vectors*. 2012;5:62.
12. Mazzariol S, Cassini R, Voltan L et al. Heartworm (*Dirofilaria immitis*) infection in a leopard (*Panthera pardus pardus*) housed in a zoological park in north-eastern Italy. *Parasit Vectors*. 2010;3(1):25.
13. Furtado AP, Do Carmo ES, Giese EG et al. Detection of dog filariasis in Marajo Island, Brazil by classical and molecular methods. *Parasitol Res*. 2009;105(6):1509-15.
14. Dyab AK, Galal LA, Mahmoud Ael-S et al. Xenomonitoring of different filarial nematodes using single and multiplex PCR in mosquitoes from Assiut Governorate, Egypt. *Korean J Parasitol*. 2015;53(1):77-83.
15. Méndez JC, Carretón E, Martínez-Subiela S et al. Acute phase protein response in heartworm-infected dogs after adulticide treatment. *Vet Parasitol*. 2015;209(3-4):197-201.
16. Hwang IK, Lee HY, Kim MH et al. Development of real-time PCR assay for genetic identification of the mottled skate, *Beringraja pulchra*. *Forensic Sci Int*. 2015;255:80-4.
17. Carracedo A, Bär W, Lincoln P et al. DNA commission of the international society for forensic genetics: guidelines for mitochondrial DNA typing. *Forensic Sci Int*. 2000;110(2):79-85.
18. Dawnay N, Ogden R, McEwing R et al. Validation of the barcoding gene COI for use in forensic genetic species identification. *Forensic Sci Int*. 2007;173(1):1-6.
19. Hebert PD, Ratnasingham S, deWaard JR. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proc Biol Sci*. 2003;270 Suppl 1:S96-9.
20. Hebert PD, Cywinska A, Ball SL et al. Biological identifications through DNA barcodes. *Proc Biol Sci*. 2003;270(1512):313-21.