



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Short Communication

In Vitro Antileishmanial Effects of Saffron Compounds, Crocin and Stigmasterol, on Iranian Strain of *Leishmania major* (MHOM/IR/75/ER)

*Reza RANJBAR¹, Peyman SHAYANFAR¹, Mahmood MANIATI²

1. Molecular Biology Research Center, Systems Biology and Poisonings Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran
2. English Department, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Received 15 May 2019
Accepted 10 Aug 2019

Keywords:
Crocin;
Stigmasterol;
Leishmania major;
Promastigote;
Amastigote;
In vitro

***Correspondence**
Email:
ranjbarre@gmail.com

Abstract

Background: Due to numerous side effects of common drugs in treatment of leishmaniasis, new therapeutic approaches focus on herbal compounds. Therefore, we aimed to determine the effect of crocin and stigmasterol on in-vitro growth of promastigotes and amastigotes of *Leishmania major* in the Department of Parasitology, Pasteur Institute, Tehran, Iran in 2018.

Methods: The effect of different concentrations of crocin and stigmasterol were evaluated by determining their in-vitro inhibitory effects on promastigotes and amastigotes of the *L. major* using MTT assay.

Results: The fatality rate was 65.27% and 71.96% for crocin and stigmasterol respectively at 24 h post-culture in concentration of 50 µg/mL. The mean inhibitory effect of crocin and stigmasterol on *L. major* amastigotes after 72 h were 52.22% and 38.96%.

Conclusion: The crocin and stigmasterol had efficient adverse effects on promastigote and amastigotes of *L. major*, hence, further studies on the anti-leishmanial effects of these herbal compounds in human and animal models are recommended.

Introduction

Leishmaniasis is a group of protozoan diseases commonly found in both human and animal (Zoonoses) (1). Cutaneous

leishmaniasis (also known as oriental sore, tropical sore, chiclero ulcer, Aleppo boil) is the most common form of leishmaniasis in



Copyright © 2021 Ranjbar et al. Published by Tehran University of Medical Sciences
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license
(<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.

both dry (urban) and wet (rural) forms world-wide including Iran (2). It has been considered as one of the six major diseases in the World Health Organization's Tropical Diseases Survey (3), about 19000 cases of reported annually (4, 5).

Unfortunately, despite the high prevalence of cutaneous leishmaniasis in Iran, no efficient prevention and treatment method has been proposed. Therefore, efforts to discover new drugs with the least side effects need to be continued more decisively (6). Systemic therapies are the first line of treatment against leishmaniasis (7). These compounds are toxic to parasites. Long duration of treatment, parasites resistance, high expense, and severe toxicity on the heart, liver and kidney could be referred as the most notable disadvantages of these drugs. Keeping in view these facts, researchers are trying to use plant compounds to treat leishmaniasis (8). Commonly, useful botanical compounds are identified through study of plant extracts. In a study, it was collected the Iranian traditional herbal remedies against leishmaniasis. Many native plants constituents such as *Zajuria multiflora* Boiss, *Lawsonia inermis*, *Calendula officinalis*, *Nerium oleander*, etc. have been effective in controlling the disease in different studies (9).

Crocin is one of the most consequential alkaloids ingredients in plants such as saffron, and many of its helpful effects on health, such as antioxidant, anti-cancer, learning and memory invigoration effects, have been proven in many studies (10). On the other hand, stigmaterol is also an herbal sterol with beneficial therapeutic functions, including anti-inflammatory, anti-cancer effects, as well as protective influence on the immune system (11).

Considering the importance of using plant compounds, the main objective of the present study was the in-vitro investigation of beneficial effects of the two herbal compounds, crocin and stigmaterol, on *L. major* standard strain (MHOM/IR/75/ER).

Materials and Methods

Our study has the control group included *L. major* promastigotes, uninfected macrophage and infected macrophage by *L. major* without crocin and stigmaterol, and case group including promastigotes of *L. major*, uninfected macrophages and macrophage infected with *L. major* treated with different doses of crocin and stigmaterol. This project was performed in the Department of Parasitology, Pasteur Institute, Tehran, Iran in 2018.

Parasite culture

L. major standard strains (MHOM/IR/75/ER) were prepared from Parasitology Department of Pasteur Institute of Iran and cultured in flasks containing RPMI-1640 medium with 7%-10% inactivated FCS.

Preparation of different concentrations of crocin and stigmaterol

About 1 mg of crocin and stigmaterol powders purchased from Sigma Company, dissolved in 200 μ l of methanol separately and various concentrations including 1.61, 3.12, 6.24, 12.5, 25, 50 μ g / mL were prepared from both compounds dissolving in the RPMI-1640 medium and stored in the 4 °C (12). The concentrations of crocin and stigmaterol were calculated according to Table 1.

Promastigotes viability Determination

10^7 parasites were cultured in 96-well plates in RPMI-1640 medium with 10% bovine serum. Crocin and Stigmaterol, with final concentrations of 1.61, 3.12, 6.24, 12.5, 25, 50 μ g/mL were added to the wells in three repetitions (triplicate) (13). The wells were examined 24 h post-culture by MTT method in which 20 μ l of MTT solution was added into wells and incubated at 37 °C in the dark for 2 to 5 hours. After centrifuging at 2500 rpm, the supernatant was discarded and 20 μ l of DMSO added and light absorption at 650 nm was examined (14).

Table 1: Calculation of different concentrations of crocin and stigmometol compounds

Ccompounds (Concentration ($\mu\text{g}/\text{ml}$))						
Crocin						
Mass (Microgram)	1.61	3.12	6.24	12.5	25	50
Concentration (micromolar)	1.6377	3.1936	6.3872	12.7948	25.5896	51.179
Volume (Milliliter)	1					
m. weight (g/mol)	976.96					
Stigmasterol						
Mass (Microgram)	1.61	3.12	6.24	12.5	25	50
Concentration (Micromolar)	3.877	7.754	15.1203	30.2891	60.5782	121.1563
Volume (Milliliter)	1					
m. weight (g/mol)	412.69					

Amastigotes viability Determination

To evaluate the EC_{50} of crocin and stigmasterol on *L. major* amastigotes, mouse macrophage J774A.1 cells were used. Primarily, the macrophage cells cultured in RPMI-1640 and incubated at 37 °C with CO_2 5% for allowing them to adhere and were then exposed to *L. major* promastigotes. To infect macrophages, 10^6 promastigotes of the parasite added to each well-containing macrophage and kept at 37 °C with CO_2 5%. After 6 h, the supernatant discarded to remove the unshielded macrophages, fresh culture medium were replaced, and solution light absorption was measured by MTT assay.

Statistical Analysis

The experiment was arranged in Completely Randomized Design (CRD) with three replications per treatment. All data were assessed by using SPSS version 20.0. (Chicago, IL, USA). The homogeneity of variance was checked using One-Sample Kolmogorov-Smirnov Test. Differences between treatment means in all

experiments were analyzed based on Scheffea test using the probability of five percent.

Results

The mean of promastigotes counted in wells containing various concentrations of crocin and stigmasterol (0, 1.61, 3.12, 6.24, 12.5, 25, 50 $\mu\text{g}/\text{mL}$) were 15.77, 12.87, 12.61, 11.54, 8.51, 5.79, 5.49 (Fig. 1.A) and 7.12, 6.41, 5.85, 5.87, 4.55, 2.49, 1.99, respectively (Fig. 1.B). The results of One-Sample Kolmogorov-Smirnov Test clearly confirmed the homogeneity of variance in both of crocin ($P=0.79$) and stigmasterol ($P=0.52$) groups. According to the findings, the mean number of promastigotes showed highly significant differences in different concentrations of crocin ($P\leq 0.0001$). The mean number of promastigotes counted stigmasterol was also significantly different ($P\leq 0.0001$). In different concentrations ranging from 0 to 50 $\mu\text{g}/\text{mL}$.




Fig. 1: The means and standard errors of promastigotes counted in different concentrations of A) crocin, and B) stigmaterol

According to Fig. 2, the graphs determined that all examined concentrations of crocin and stigmaterol had a significant effect on the number of parasites compared to control group. In another word, by increasing the concentration, there was a significant decrease in the number of parasites. After 24 h, the concentration of crocin and 19 of stigmaterol causes 50% of the parasite population to be dead. The lowest concentration of stigmaterol showed 10.07% growth inhibitory effect after 24 h and higher concentrations were more efficient so that at 50 and 25 µg/ml, the percentage of fatality was reported to be 71.96% and 65.03%, respectively.

The light absorption indicated less parasites exist in higher concentrations

As shown in Table 1, the optical density (15) decreased when the concentration of extractions increased from 0 to 50 µg/mL. The OD was 0.95 in 1.61 µg/mL of Crocin; meanwhile, the OD reached the lowest number (0.49) in the highest concentration (50 µg/mL). The same pattern could observe in various concentrations of stigmaterol showing that more the concentrations increased, more the OD decreased (Table 2).

Table 2: The measured OD of crocin and stigmaterol in various concentrations

Concentration (µM)		OD*	
Crocine	Stigmaterol	Crocine	Stigmaterol
0 **	0 **	1.40 ±0.003	9.80 ±0.08
1.6	3.8	0.95 ±0.07	0.68 ±0.11
3.1	7.7	0.81 ±0.01	0.63 ±0.08
6.3	15.1	0.85 ±0.02	0.61 ±0.44
12.7	30.2	0.77 ±0.02	0.49 ±0.13
25.5	60.5	0.71 ±0.04	0.32 ±0.02
51.1	121.1	0.49 ±0.02	0.22 ±0.15

* Data are presented in Mean ± SD

** Control group

L. major promastigotes exposing to high concentrations has been led to death more in comparison to counterparts exposed to low concentrations in both crocin and stigmasterol extractions (Fig. 2). The crocin had better in-

hibitory effect in lower concentrations. However, in higher concentrations including 25 $\mu\text{g}/\text{mL}$ and 50 $\mu\text{g}/\text{mL}$, the stigmasterol extraction had efficacy compared to crocin (Fig. 3).




Fig. 2: The growth inhibitory effect of different concentrations of crocin and stigmasterol on *L. major* promastigotes Standard strain in RPMI 1640 after 24 h




Fig. 3: The crocin and stigmasterol inhibitory effect on *L. major* promastigote standard strain in different concentrations ranging from 0 to 50 $\mu\text{g}/\text{mL}$

The mean of amastigotes counted in each macrophage decreased after 72 h in control, Crocin-treated and Stigmasterol-treated groups in three treatments. Table 3 shows the mean number of amastigotes counted in 100 macrophages in control, Crocin-treated and Stigmasterol-treated groups in the three treat-

ments (Table 2). The mean inhibitory effect of these compounds on *L. major* amastigotes has been measured 52.2% and 38.96% for crocin and Stigmasterol, respectively. The results indicate the higher efficiency of crocin anti-leishmaniasis effect compared to stigmasterol (Table 2).

Table 3: The crocin and stigmaterol EC50 inhibition activity on *L. major* promastigote standard strain

Groups	Mean (100 macrophages $\times 10^6$)		Growth inhibitory (%)	
	Crocin	Stigmaterol	Crocin	Stigmaterol
Treated Control 1	2.25	2.79	0	0
Crocin/ stigmaterol treated 1	1.06	1.61	52.48	42.2
Treated Control 2	2.17	3.04	0	0
Crocin/ stigmaterol treated 2	0.99	1.88	57.64	38.18
Treated Control 3	2.21	2.89	0	0
Crocin/ stigmaterol treated 3	1.18	1.83	46.48	36.51

Discussion

Leishmaniasis is a group of parasitic diseases caused by the protozoan *Leishmania* species (16). Annually, between 1.5 and 2 million people suffer from cutaneous leishmaniasis globally, 90% of whom live in 9 countries, including Iran (17). Since the efforts for designing vaccine has failed in clinical trials so far, there is a dire need for new treatments. Herbal bioactive compounds extracted from a variety of plants exhibit anti-leishmanial properties (18).

Traditional medicine, particularly those are based on the use of plants, has been regarded as an ancient root stone to prevent and treat many infections from centuries ago to now (19, 20). Natural products and their derivatives that are originated from plants have potential activities against microbial agents including parasites (21, 22).

Stigmaterol has been shown to stimulate cytokines release by which the differentiation of CD4+ T cells into Th1 cells occurs (23). Macrophages, are known as parasite host cells and presents parasitic antigens to immune cells (24). Stigmaterol can stimulate cellular immunity and activates macrophage cells and natural killers cells (NKs) (25).

In a study, anticoagulant activity and the effects of immune modulation of tannins were investigated. The effects of polyphenol on intracellular leishmaniasis parasites are due to

activation of macrophages instead of direct antiparasitic activity (26).

Stigmaterol induce the caspase-3 activation and anti-apoptotic protein Bcl-2 decrement and thus results in apoptosis (27). *L. major* contributes to its survival in the host body by preventing the release of cytochrome C from mitochondria and as a result, harnesses Caspase-3 activation and apoptosis in infected macrophage (28). Phytosterols such as stigmaterol trigger the activation of Caspase-3 and induce internal pathways apoptosis (mitochondrial) (29). In addition to activating caspase, stigmaterol increases the production and activation of certain enzymes involved in the *L.* delimitation, such as calmodulin and cellular kinases (30).

The investigation of Inhibitory effect of saffron and its major components, Safranal and Crocin, revealed that *Helicobacter pylori* was vulnerable to saffron aquatic and alcoholic extracts (31). Safranal and crocin had more antibacterial effects in comparison to alcoholic extracts (15). In this regard, the Iranian native plants had potential therapeutic effects on cutaneous leishmaniasis. They pointed to medicinal plants of different parts in Iran, although the saffron compounds were not named due to expensive, the high prevalence of leishmaniasis in Khorasan Province and the abundance of this product in the city shows the importance of the effective compounds in this Area (32).

Two herbal compounds of saffron, stigmasterol and crocin had concentration-dependent inhibitory effect on the growth of *L. major* amastigotes and promastigotes in macrophage cells. As expected, we deduced that the higher concentration of stigmasterol and crocin accompany the lower growth rate of the parasite. Therefore, the survival rate of the parasite promastigote and amastigote is also dependent on the concentration of these compounds.

Conclusion

Since stigmasterol and crocin have anti-leishmaniasis influence in-vitro, the in-vivo studies in animal models are recommended in achieving the appropriate medicine combination in the treatment of cutaneous leishmaniasis.

Acknowledgements

This study was financially supported in part by “Clinical Research Development Center of Baqiyatallah Hospital.

Conflict of interest

The authors declare that there is no conflict of interests.

References

1. Reithinger R, Dujardin J-C, Louzir H, et al. Cutaneous leishmaniasis. *Lancet Infect Dis.* 2007;7(9):581-96.
2. Schwartz E, Hatz C, Blum J. New world cutaneous leishmaniasis in travellers. *Lancet Infect Dis.* 2006;6(6):342-9.
3. Alvar J, Vélez InD, Bern C, et al. Leishmaniasis worldwide and global estimates of its incidence. *PLoS One.* 2012;7(5):e35671.
4. Liu D, Uzonna JE. The early interaction of *Leishmania* with macrophages and dendritic cells and its influence on the host immune response. *Front Cell Infect Microbiol.* 2012;2:83.
5. Mehravaran A, Jaafari MR, Jalali SA, et al. The role of ISCOMATRIX bilayer composition to induce a cell mediated immunity and protection against leishmaniasis in BALB/c mice. *Iran J Basic Med Sci.* 2016;19(2):178-86.
6. Farahmand M, Nahrevanian H, Shirazi HA, et al. An overview of a diagnostic and epidemiologic reappraisal of cutaneous leishmaniasis in Iran. *Braz J Infect Dis.* 2011;15(1):17-21.
7. Pourmohammadi B, Motazedian M, Handjani F, et al. Glucantime efficacy in the treatment of zoonotic cutaneous leishmaniasis. *Southeast Asian J Trop Med Public Health.* 2011;42(3):502-8.
8. Sadeghian G, Ziaei H, Bidabadi LS, et al. Decreased effect of glucantime in cutaneous leishmaniasis complicated with secondary bacterial infection. *Indian J Dermatol.* 2011;56(1):37-9.
9. Bahmani M, Saki K, Ezatpour B, et al. Leishmaniasis phytotherapy: Review of plants used in Iranian traditional medicine on leishmaniasis. *Asian Pac J Trop Biomed.* 2015;5(9):695-701.
10. Rezaee R, Mahmoudi M, Abnous K, et al. Cytotoxic effects of crocin on MOLT-4 human leukemia cells. *J Complement Integr Med.* 2013;10(1). doi: 10.1515/jcim-2013-0011.
11. Kaur N, Chaudhary J, Jain A, et al. Stigmasterol: a comprehensive review. *Int J Pharm Sci Res.* 2011;2(9):2259-2265.
12. Ghaffarifar F. *Leishmania major*: in vitro and in vivo anti-leishmanial effect of cantharidin. *Exp Parasitol.* 2010;126(2):126-9.
13. Khademvatan S, Gharavi MJ, Rahim F, et al. Miltefosine-induced apoptotic cell death on *Leishmania major* and *L. tropica* strains. *Korean J Parasitol.* 2011;49(1):17-23.
14. Maroufi Y, Ghaffarifar F, Dalimi A, et al. A study on the cytotoxic effect of cantharidin on *Leishmania major* promastigote and amastigote survival in vitro. *KAUMS Journal(FEYZ).* 2012;16(5):406-413.
15. Nakhaei M, Khaje-Karamoddin M, Ramezani M. Inhibition of *Helicobacter pylori* growth in vitro by saffron (*Crocus sativus* L.). *Iran J Basic Med Sci.* 2008;11:91-6.

16. Kaye P, Scott P. Leishmaniasis: complexity at the host-pathogen interface. *Nat Rev Microbiol.*2011;9(8):604-15.
17. Monge-Maillo Ba, López-Vélez R. Therapeutic options for old world cutaneous leishmaniasis and new world cutaneous and mucocutaneous leishmaniasis. *Drugs.*2013;73(17):1889-920.
18. De Monte C, Bizzarri B, Gidaro MC, et al. Bioactive compounds of *Crocus sativus* L. and their semi-synthetic derivatives as promising anti-*Helicobacter pylori*, anti-malarial and anti-leishmanial agents. *J Enzyme Inhib Med Chem.*2015;30(6):1027-33.
19. Khomarlou N, Aberoomand-Azar P, Lashgari AP, et al. Essential oil composition and in vitro antibacterial activity of *Chenopodium album* subsp. striatum. *Acta Biol Hung.* 2018; 69(2): 144-55.
20. Mirzaie A, Halaji M, Dehkordi FS, et al. A narrative literature review on traditional medicine options for treatment of corona virus disease 2019 (COVID-19). *Complement Ther Clin Pract.* 2020;40:101214
21. Jonaidi Jafari N, Kargozari M, Ranjbar R, et al. The effect of chitosan coating incorporated with ethanolic extract of propolis on the quality of refrigerated chicken fillet. *Journal of Food Processing and Preservation.* 2018; 42(1): e13336.
22. Aminnezhad S, Kermanshahi RK, Ranjbar R. Evaluation of synergistic interactions between cell-free supernatant of *Lactobacillus* strains and Amikacin and genetamicin against *Pseudomonas aeruginosa*. *Jundishapur J Microbiol.* 2015;8(4):e16592.
23. Nylén S, Gautam S. Immunological perspectives of leishmaniasis. *J Glob Infect Dis.*2010;2(2):135-46.
24. Huang C-H, Cheng J-Y, Deng M-C, et al. Prebiotic effect of diosgenin, an immunoactive steroidal sapogenin of the Chinese yam. *Food Chem.*2012; 132(1):428-32.
25. Dar NJ, Hamid A, Ahmad M. Pharmacologic overview of *Withania somnifera*, the Indian Ginseng. *Cell Mol Life Sci.*2015;72(23):4445-60.
26. Kolodziej H, Kiderlen AF. Antileishmanial activity and immune modulatory effects of tannins and related compounds on *Leishmania* parasitised RAW 264.7 cells. *Phytochemistry.* 2005;66(17):2056-71.
27. Kabeer FA, Sreedevi GB, Nair MS, et al. Isoledeoxyelephantopin from *Elephantopus scaber* (Didancao) induces cell cycle arrest and caspase-3-mediated apoptosis in breast carcinoma T47D cells and lung carcinoma A549 cells. *Chin Med.*2014;9:14.
28. Looi CY. Pharmacological Activities and Chemical Constituents of. *J Med Sci.*13(4):236-43.
29. Sirisha N, Sreenivasulu M, Sangeeta K, et al. Antioxidant properties of *Ficus* species—a review. *Int J Pharmtech Res.*2010;2.
30. Brasili E, Praticò G, Marini F, et al. A non-targeted metabolomics approach to evaluate the effects of biomass growth and chitosan elicitation on primary and secondary metabolism of *Hypericum perforatum* in vitro roots. *Metabolomics.*2014;10:1186-1196.
31. Rosenthal E, Marty P. Recent understanding in the treatment of visceral leishmaniasis. *J. Postgrad Med.* 2003;49(1):61-8.
32. Moghaddas E, Khamesipour A, Mohebbali M, et al. Iranian Native Plants on Treatment of Cutaneous Leishmaniasis: A Narrative Review. *Iran J Parasitol.* 2017;12(3):312-322.