



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Short Communication

Intestinal Parasitic Infections in People Referring to the Central Laboratory of Meshkin Shahr County, Ardabil Province, Iran

Pooria Asadi ¹, Zabihollah Zarei ¹, Mehdi Mohebali ¹, Zahra Alizadeh ¹, Faezeh Najafi ¹,
*Shahrokh Izadi ¹, *Zahra Heidari ^{2,3}

1. Department of Medical Parasitology and Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
2. Zoonoses Research Center (ZRC), Ardabil University of Medical Sciences, Ardabil, Iran
3. Department of Medical Microbiology, Parasitology and Immunology, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran

Received 23 Sep 2023
Accepted 04 Nov 2023

Keywords:

Intestinal parasitic infections;
Prevalence;
Protozoans;
Helminths;
Iran

***Correspondence**

Email:
za.heidari@arums.ac.ir

Abstract

Background: Intestinal parasitic infections are still a considerable global public health problem. We aimed to determine the frequency of intestinal parasitic infections among people referring to the central laboratory of Meshkin Shahr City, Ardabil Province, Iran.

Methods: In this cross-sectional survey, 460 fecal samples were collected randomly from persons referred to the central laboratory of Meshkin Shahr City, from January to June 2022. The samples were examined by direct wet-mount, Trichrome and modified Ziehl-Neelsen staining, formalin ethyl acetate sedimentation, and agar plate culture.

Results: The frequency of intestinal parasites was 15.7% (72 out of 460 cases), with some people with numerous intestinal parasites. The frequency of protozoan infections (13.9%) was higher than the helminthic infections (2.6%). *Blastocystis* spp. (8.1%) was the most prevalent detected intestinal protozoan. *Entamoeba coli* (5.7%), *Dicrocoelium dendriticum* (2.2%), *Giardia lamblia* (1.5%), *Fasciola* spp. (0.2%), and *Hymenolepis nana* (0.2%) were other detected parasites.

Conclusion: In spite of betterment of the health condition in Iran and reduction of parasitic infection, intestinal parasitic infections are still a considerable public health issue in some parts of Iran.



Introduction

Intestinal parasitic infection (IPIs) caused by protozoa and soil-transmitted helminths is brought up one of the significant public health issues due to its high prevalence and distribution throughout the world, especially in the populations of tropical and subtropical areas of developing countries (1). Several species of STH, *Ascaris lumbricoides*, *Trichuris trichiura*, and hookworms (*Ancylostoma duodenale* and *Necator americanus*) included as part of the neglected tropical diseases (NTDs) lately updated by the WHO (2, 3). It is approximated more than one-third of the world's people are infected with intestinal parasites, which constitutes a global health burden that causes clinical complications and mortality (4, 5). The IPIs cause the yearly loss of about 39 million disability-adjusted life years, resulting in widespread health and economic problems (6, 7). Intestinal parasites can cause a broad range of symptoms, from asymptomatic carriage, which is observed in about 90% of infected people, to diarrhea, abdominal pain, intestinal bleeding, anemia, malabsorption, malnutrition, general malaise and weakness that may affect learning capacities and physical growth (8, 9).

Indirect transmission of intestinal parasites usually occurs through ingesting contaminated water, food, and soil (fecal-oral). Some species have the possibility of direct transmission through person-to-person or animal-to-human contact (10, 11).

Several environmental, geographic, and social factors cause differences in the prevalence and distribution of IPIs from one region to another. The prevalence of intestinal parasites is closely related to socio-economic status, educational levels, environmental factors, insufficient medical care, lack or inadequate access to safe drinking water sources, improper hygiene, poor health practices and education (11, 12). In Iran, a wide range of intestinal parasite prevalence between 4.7 to 56% has been reported in apparently healthy individu-

als (13). However, some parasitic infections such as hookworms and ascariasis have dropped significantly in the last years (13, 14). Nevertheless, giardiasis is the most common protozoan infection in Iran (15).

So far, no study on the frequency of intestinal parasitic infections in Meshkin Shahr City. We aimed to determine the frequency of intestinal parasites among people referring to the Meshkin Shahr Central Laboratory.

Methods

Study area

Meshkin Shahr County, with an area surface of about 3,500 km², is placed in Ardabil Province in Iran (38°26'N 47°45'E). The County is divided into four sections: Meshkin-e Sharqi District, the Central District, Moradlu District and Arshaq District and also It has six cities: Meshkin Shahr, Qasabeh, Lahrud, Moradlu, Razi and Fakhrabad. According to the 2016 census, it has inhabitants of about 149,941 with 45,999 households. Meshkin Shahr County is approximately 1400 meters above sea level, and Sabalan Mountain has located 25 km from this County. It has four climates: hot and dry Mediterranean, temperate Mediterranean, cold steppe and Cold Mountain. The length of the dry and semi-dry, and frosty season is five to eight months, and the average annual rainfall is 300 mm. Most of its vegetation consists of steppes, artichokes, hornbeam, oaks, and horrida, which provide enormous pastures for livestock, a good opportunity for the transmission and spread of zoonotic diseases in the region.

Study design

The required sample size was calculated as 202 concerning the frequency of parasitic infections in the region. To increase the studies accuracy by applying the design effect, 460 participants were included.

This cross-sectional study was performed in Meshkin Shahr City for six months between January and June 2022. Four hundred and sixty (460) fecal samples (from 284 males and 176 females) were collected by simple random sampling among individuals referring to the central laboratory of the city. At first, a screw-capped plastic container was given to each participant; then, during sampling, a standard predesigned questionnaire including gender, age, education level, occupation, type of residency and clinical indications and symptoms (e.g., nausea, diarrhea, abdominal pain and cramp) was completed through interviews by a trained clinical laboratory expert. The collected containers in insulated boxes were transported to the health research station of Meshkin Shahr, Tehran University of Medical Sciences. Finally, the samples were also processed on the same day of collection.

Stool examinations

All fresh fecal specimens were macroscopically evaluated for consistency, color, the presence of blood, mucus, and adult intestinal worms. The samples were examined using the direct wet-mount and formalin ethyl acetate sedimentation techniques to diagnosis the presence of any parasite. Subsequently, a part of each collected sample (approximately 2 g) was cultured on agar plates, and each plate was observed by stereomicroscope to detect *Strongyloides stercoralis* and *Hookworm*. The trichrome stain was applied for the detection of protozoan infections. A thin smear was prepared from diarrhea specimens, and further evaluation was performed using a modified Ziehl-Neelsen staining technique to detect coccidian parasites. Eventually, all the slides were observed under a light microscope for at least 20 min using 100×, 400× magnification and the stained slides with 1000× as final magnification.

Ethics Statement

First, the protocol of this cross-sectional study was evaluated and approved by the Eth-

ics Committee of Tehran University of Medical Sciences (ethical code: 400000707) following the Helsinki Declaration and Guidelines. Information about the aim of the study was provided to the adult individuals and parents/legal guardians of the children; also, written informed consent was obtained from all individuals.

Statistical analysis

Data processing and analysis were conducted on SPSS 26 software (IBM Corp., Armonk, NY, USA). The chi-square test was applied to analyze the association between variables, and also Significance level was accepted at P values < 0.05 .

Results

Over six months, 460 stool specimens from the Meshkin Shahr residents, along with 284 (61.7%) men and 176(38.3%) women, were gathered. The age of the people was between 1 year as the youngest participant to 86 years as the oldest person studied, with an average age of 31.23 ± 34.58 years, where most of the participants were middle-aged and in the age group of 30-39 (25.4%). Only 13.7% (63/460) of the participants had academic (university) education, and 22.8% (105/460) of them were illiterate. Two hundred twenty-four individuals (48.7%) lived in the rural regions and two hundred thirty-six individuals (51.3%) lived in urban regions.

The frequency of detected intestinal parasites in the participants is provided in Table 1. At least one species of intestinal parasite was found in 72 (15.7%) participants.

The frequency of infection in males and females was (16.9%) and (13.6%) respectively. Intestinal parasitic infections showed no significant correlation with gender ($P= 0.34$). The maximum rate of infection was seen in the 60-69 age group (26.1%), but no significant association was reported between the age group and the frequency of infection ($P= 0.30$). The current survey demonstrates a significant as-

sociation between clinical manifestations and infection with intestinal parasites ($P= 0.001$). However, occupation, education, and place of

residence were not associated with the parasitic intestinal infection rates.

Table 1: Frequency of IPIs according to parasite species and number of infections among the participants (n=460)

Variable	Parasite's species	Number of infected	Frequency (%)
Single	<i>Blastocystis</i> spp.	29	6.3
	<i>Entamoeba coli</i>	21	4.6
	<i>Giardia lamblia</i>	5	1.1
	<i>Dicrocoelium dendriticum</i>	6	1.3
	<i>Fasciola</i> spp.	1	0.2
	<i>Hymenolepis nana</i>	1	0.2
Double	<i>Blastocystis</i> spp., <i>Dicrocoelium dendriticum</i>	3	0.7
	<i>Blastocystis</i> spp., <i>Entamoeba coli</i>	3	0.7
	<i>Blastocystis</i> spp., <i>Giardia lamblia</i>	1	0.2
	<i>Entamoeba coli</i> , <i>Dicrocoelium dendriticum</i>	1	0.2
Triple	<i>Blastocystis</i> spp., <i>Giardia lamblia</i> , <i>Entamoeba coli</i>	1	0.2
	Total	72	15.7

Discussion

In this cross-sectional study, 15.7% of participants were positive for intestinal parasites. According to our knowledge, this research was the first attempt to illustrate the frequency of IPIs in this area. This finding was similar to a study performed in Tehran in which the prevalence of IPIs was reported as 17% among patients visiting a referral hospital (16). In addition, the prevalence of the intestinal parasite in people referred to different medical laboratories in Babol was 12.1% (17, 18). A nationwide study of the public population reported a prevalence rate of 19.3% for IPIs in Iran (19). In the earlier studies performed on specific community groups in Ardabil, near Meshkin Shahr, the prevalence of IPIs was reported to be 3.1%, 10%, and 4.5% in the applicants receiving health card, patients with malignancy, patients referred to laboratories of two hospitals affiliated with Ardabil university of medical sciences, respectively (20, 21). The prevalence found in our study is higher than previous reports from the south of Tehran, Karaj,

and Qazvin, with IPIs prevalence of 10.7%, 4.7%, and 5.8%, respectively (22-24). However, some studies reported a remarkable prevalence of 32.2% in patients with gastrointestinal diseases in Nahavand County, 32.7%, and 34.2% in inhabitants of Roudehen and Rudbar-e Jonub (13, 25, 26).

Our findings and other studies indicated that despite advancements in hygiene and public/personal health measures, there are still insufficient performances for controlling intestinal parasites, and IPIs are still a significant public health issue in Iran and developed countries (13, 17). Several factors, such as weather conditions, agriculture, and access to health facilities, study period, lifestyle, and the laboratory techniques employed for stool examination, differences in study populations, and sanitary conditions in the investigated areas, were probable causes of the differences between our findings and those of the previous studies completed in Iran in terms of the general prevalence of IPIs.

In the current study, the frequency of protozoa infections was higher at 13.9% than that

of helminth infections at 2.6%. The high frequency of protozoan parasites is likely due to their high reproductive rate and easy route of infection via contaminated water and foods (27). In addition, several studies conducted in different parts of Iran indicate that in recent years, the prevalence of intestinal helminth infections in Iran has decreased; this may be due to overall improvements in sanitation in Iran (28, 29). Our results show that about 2% of study participants were infected with multiple parasites at the same time. *Blastocystis* spp. was the most common intestinal parasite in co-infections. The prevalence of mixed infection differs in different studies conducted around the world (30). This may be due to the same transmission route in many protozoan species. *Blastocystis* spp., *E. coli*, and *G. lamblia* were the most common intestinal protozoa among participants in the current study, consistent with other studies conducted in Iran (31, 32).

In the present study, the most common intestinal parasite was *Blastocystis* spp. 8.1%, similar to other surveys conducted in Iran and worldwide (13, 33). Studies on apparently healthy population groups in Iran conducted over the past few decades have reported the variable prevalence of *Blastocystis* infection to be between 7.5 to 28.4% (26, 34).

According to our results, the frequency of *E. coli* was 5.7%. Based on the studies conducted in Iran, the prevalence of *E. coli* was reported up to 18.9% (35). As some protozoans, such as *Blastocystis* spp. and *E. coli*, are transmitted to humans through the fecal-oral route, their presence is a suitable indicator of poor hygiene, close animal contact, and contaminated food or water consumption (13, 26).

G. lamblia infection is observed worldwide, and this parasite is one of the most common protozoa that can cause diarrhea (13). The frequency of *G. lamblia* in these study participants was 1.5%. However, based on the laboratory results of several studies conducted in the different regions of Iran, the prevalence of

Giardia has decreased from 33.9% to 1.2% in recent years compared to the previous decades (26, 34).

In terms of helminth infection, three types of parasite ova, including *D. dendriticum*, *H. nana*, and *Fasciola* spp., were found in the infected samples. The prevalence of helminth infections in Iran has decreased dramatically in recent years, while some helminths with direct fecal-oral transmission, such as *H. nana* and *Enterobius vermicularis*, remain common in some parts of the country (36). Despite the human cases of dicrocoeliasis reported in the northern regions of Iran, the *D. dendriticum* infections reported in the current study were pseudo infections due to consuming contaminated livestock liver (16). The low rate of intestinal helminth infections in Meshkin Shahr is consistent with the findings of recent reports from other regions of Iran (16, 25).

The current study examined several possible risk factors correlating with intestinal parasitic infections and found that clinical symptoms were significantly associated with intestinal parasitic infections ($P < 0.05$), consistent with the results of previous studies (18, 25).

In the current study, the frequency of IPIs was slightly lower in females (13.6%) than in males (16.9%). Although intestinal parasite infection showed no significant difference with sex, indicating equal exposure to infectious agents; similar results have also been reported in other surveys (37).

Based on our findings, the frequency of IPIs in the age group 60-69 years was 26.1% higher than in other age groups, possibly due to the low personal hygiene and education level of older people and lack of awareness of the transmission of IPIs. Nevertheless, there was no significant correlation between the frequency of IPIs and age groups, consistent with other reported studies (13, 16).

The frequency of intestinal parasites showed no significant correlation with educational level or occupation, although the highest fre-

quency was found among retired participants, 22.2% (16, 18).

There was no significant correlation between the infection and residential status, which is similar to the findings reported previously (38).

The present study was subjected to some limitations. The study was non-blinded. As the collection period was short. Another limitation was that due to facilities and financial constraints, we had not applied all the laboratory techniques for detection of parasites.

Conclusion

Although in recent years, due to the improvement of the health situation in Iran, the prevalence of IPIs in Iran has decreased. IPIs are still an important public health problem in Iran. As most of the detected parasites are directly transmitted, raising public awareness, health education, infected individuals follow-up, and treatment should be considered to prevent this infection. In this context, the results of the current study can be used as a basis for the development of public health strategies and prevention programs for IPIs.

Acknowledgements

The authors are thankful to all participants in this study. This work was financially supported by Zoonoses Research Center, Ardabil University of Medical Sciences and Project No: 400000707.

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Brooker S, Clements AC, Bundy DA. Global epidemiology, ecology and control of soil-transmitted helminth infections. *Adv Parasitol.* 2006;62:221-61.
2. WHO. Neglected tropical diseases. 2017. Available online at: <https://www.who.int/news-room/questions-and-answers/item/neglected-tropical-diseases> (accessed May 4, 2021). 2018.
3. Periago MV, García R, Astudillo OG, et al. Prevalence of intestinal parasites and the absence of soil-transmitted helminths in Añatuya, Santiago del Estero, Argentina. *Parasit Vectors.* 2018;11(1):638.
4. Tefera T, Mebric G. Prevalence and predictors of intestinal parasites among food handlers in Yebu town, southwest Ethiopia. *PLoS One.* 2014;9(10):e110621.
5. Chelkeba L, Mekonnen Z, Alemu Y, Emanu D. Epidemiology of intestinal parasitic infections in preschool and school-aged Ethiopian children: a systematic review and meta-analysis. *BMC Public Health.* 2020;20(1):117.
6. Ramana K. Intestinal parasitic infections: An overview. *Ann Trop Med Public Health.* 2012;5(4):279-281.
7. Eyayu T, Kiros T, Workineh L, et al. Prevalence of intestinal parasitic infections and associated factors among patients attending at Sanja Primary Hospital, Northwest Ethiopia: An institutional-based cross-sectional study. *PLoS One.* 2021;16(2):e0247075.
8. Ahmed A, Al-Mekhlafi HM, Al-Adhroey AH, et al. The nutritional impacts of soil-transmitted helminths infections among Orang Asli schoolchildren in rural Malaysia. *Parasit Vectors.* 2012;5:119.
9. Habib A, Andrianonimiadana L, Rakotondrainipiana M, et al. High prevalence of intestinal parasite infestations among stunted and control children aged 2 to 5 years old in two neighborhoods of Antananarivo, Madagascar. *PLoS Negl Trop Dis.* 2021;15(4):e0009333.
10. Seguí R, Muñoz-Antoli C, Klisiowicz DR, et al. Prevalence of intestinal parasites, with emphasis on the molecular epidemiology of *Giardia duodenalis* and *Blastocystis* sp., in the Paranaguá Bay, Brazil: a community survey. *Parasit Vectors.* 2018;11(1):490.
11. Younes N, Behnke JM, Ismail A, Abu-Madi MA. Socio-demographic influences on the

- prevalence of intestinal parasitic infections among workers in Qatar. *Parasit Vectors*. 2021;14(1):63.
12. Sahimin N, Lim YA, Ariffin F, et al. Migrant workers in Malaysia: current implications of sociodemographic and environmental characteristics in the transmission of intestinal parasitic infections. *PLoS Negl Trop Dis*. 2016;10(11):e0005110.
 13. Abbaszadeh Afshar MJ, Barkhori Mehni M, Rezaeian M, et al. Prevalence and associated risk factors of human intestinal parasitic infections: a population-based study in the southeast of Kerman province, southeastern Iran. *BMC Infect Dis*. 2020;20(1):12.
 14. Askarian M, Ghanaie RM, Karimi A, Habibzadeh F. Infectious diseases in Iran: a bird's eye view. *Clin Microbiol Infect*. 2012;18(11):1081-8.
 15. Hatam-Nahavandi K, Mohebbali M, Mahvi AH, et al. Subtype analysis of *Giardia duodenalis* isolates from municipal and domestic raw wastewaters in Iran. *Environ Sci Pollut Res Int*. 2017;24(14):12740-12747.
 16. Zarei A, Mohebbali M, Agholi M, et al. Prevalence and Associated Risk Factors of Intestinal Parasitic Infections among Patients Visiting a Referral Hospital in Tehran Province, Iran. *Iran J Parasitol*. 2022;17(3):385-92.
 17. Badparva E, Kheirandish F, Ebrahimzade F. Prevalence of intestinal parasites in Lorestan Province, West of Iran. *Asian Pac J Trop Dis*. 2014;4:S728-S32.
 18. Beiromvand M, Panabad E, Rafiei A. Status of intestinal parasitic infections among rural and urban populations, southwestern Iran. *Asian Pac J Trop Dis*. 2019;12(3):130-6.
 19. Sayyari A, Imanzadeh F, Bagheri Yazdi S, et al. Prevalence of intestinal parasitic infections in the Islamic Republic of Iran. *East Mediterr Health J*. 2005; 11 (3):377-83.
 20. Babaei Pouya N, Razmjou E. Epidemiology of intestinal parasites among applicants receiving Health Card of Ardabil City in 2014. *J Health*. 2018;9(1):115-23.
 21. Mohammadi-Ghalehbin B, Pezeshki A, Kohansal MH, Esmailnezhad G. Frequency of intestinal parasites in patients with malignancy in Ardabil province, northwest Iran. *J Hum Environ Health Promot*. 2017;2(2):118-24.
 22. Arani AS, Alaghebandan R, Akhlaghi L, et al. Prevalence of intestinal parasites in a population in south of Tehran, Iran. *Rev Inst Med Trop Sao Paulo*. 2008;50(3):145-9.
 23. Nasiri V, Esmailnia K, Karim G, et al. Intestinal parasitic infections among inhabitants of Karaj City, Tehran province, Iran in 2006-2008. *Korean J Parasitol*. 2009;47(3):265-68.
 24. Sadeghi H, Borji H. A survey of intestinal parasites in a population in Qazvin, north of Iran. *Asian Pac J Trop Dis*. 2015;5(3):231-3.
 25. Kiani H, Haghighi A, Rostami A, et al. Prevalence, risk factors and symptoms associated to intestinal parasite infections among patients with gastrointestinal disorders in Nahavand, Western Iran. *Rev Inst Med Trop Sao Paulo*. 2016;58:42.
 26. Hemmati N, Razmjou E, Hashemi-Hafshejani S, et al. Prevalence and risk factors of human intestinal parasites in Roudehen, Tehran province, Iran. *Iran J Parasitol*. 2017;12(3):364-73.
 27. Abera B, Biadegelgen F, Bezabih B. Prevalence of *Salmonella typhi* and intestinal parasites among food handlers in Bahir Dar Town, Northwest Ethiopia. *Ethiop J Health Dev*. 2010;24(1):46-50.
 28. Sarkari B, Hosseini G, Motazedian MH, et al. Prevalence and risk factors of intestinal protozoan infections: a population-based study in rural areas of Boyer-Ahmad district, Southwestern Iran. *BMC Infect Dis*. 2016;16(1):703.
 29. Sharifdini M, Heidari Z, Hesari Z, et al. Molecular phylogenetics of *Trichostrongylus* species (Nematoda: Trichostrongylidae) from humans of Mazandaran province, Iran. *Korean J Parasitol*. 2017;55(3):279-85.
 30. Al-Delaimy AK, Al-Mekhlafi HM, Nasr NA, et al. Epidemiology of intestinal polyparasitism among Orang Asli school children in rural Malaysia. *PLoS Negl Trop Dis*. 2014;8(8):e3074.
 31. Abbaszadeh Afshar MJ, Mohebbali M, Mohtasebi S, et al. Intestinal parasites among intellectually disabled individuals in

- Iran: a systematic review and meta-analysis. *Gut Pathog.* 2021;13(1):28.
32. Mohebbali M, Keshavarz H, Afshar MJA, et al. Spatial Distribution of Common Pathogenic Human Intestinal Protozoa in Iran: A Systematic Review. *Iran J Public Health.* 2021;50(1):69-82.
33. Kuzehkanani AB, Rezaei S, Babaei Z, et al. Enteric protozoan parasites in rural areas of Bandar-Abbas, southern Iran: comparison of past and present situation. *Iran J Public Health.* 2011;40(1):80-85.
34. Mowlavi G, MirAhmadi H, Rezaeian M, et al. Prevalence of intestinal parasites in tribal parts of Khuzestan Province during 2005-07. *Govaresh.* 2008;12(4):219-28.
35. Jafari R, Fallah M, Darani HY, et al. Prevalence of intestinal parasitic infections among rural inhabitants of Hamadan city, Iran, 2012. *Avicenna J Clin Microbiol Infect.* 2014;1(2): 21445 .
36. Teimouri A, Alimi R, Farsi S, Mikaeili F. Intestinal parasitic infections among patients referred to hospitals affiliated to Shiraz University of Medical Sciences, southern Iran: a retrospective study in pre- and post-COVID-19 pandemic. *Environ Sci Pollut Res Int.* 2022;29:36911-9.
37. Sharifdini M, Ghanbarzadeh L, Barikani A, Saraei M. Prevalence of intestinal parasites among rural inhabitants of Fouman, Guilan Province, Northern Iran with emphasis on *Strongyloides stercoralis*. *Iran J Parasitol.* 2020;15(1):91-100.
38. Bahrami F, Haghghi A, Zamini G, et al. Prevalence and associated risk factors of intestinal parasitic infections in Kurdistan Province, northwest Iran. *Cogent Med.* 2018;5(1):1503777.