



Tehran University of Medical
Sciences Publication
<http://tums.ac.ir>

Iran J Parasitol

Open access Journal at
<http://ijpa.tums.ac.ir>



Iranian Society of Parasitology
<http://isp.tums.ac.ir>

Original Article

Survival of *Trichinella spiralis* and *T. pseudospiralis* in Experimentally Infected Wild Boar Muscle Tissue under Freezing and Environmental Conditions

Clara Bessi^{1,2}, Mariano Emmanuel Ercole^{1,2}, *Fernando Adrian Fariña^{1,2},
Francisco Montalvo^{1,2}, Valeria Fassa³, Marcelo Acerbo⁴, Miriam Mabel Ribicich^{1,2},
Mariana Inés Pasqualetti^{1,2}

1. Department of Parasitology and Parasitic Diseases, Faculty of Veterinary Sciences, University of Buenos Aires, Buenos Aires, Argentina
2. Institute of Animal Production Research (INPA), Faculty of Veterinary Sciences, University of Buenos Aires, Buenos Aires, Argentina
3. Department of Genetics, Faculty of Veterinary Sciences, University of Buenos Aires, Buenos Aires, Argentina
4. Department of Swine Production, Faculty of Veterinary Sciences, University of Buenos Aires, Buenos Aires, Argentina

Received 10 Feb 2024
Accepted 18 Apr 2024

Keywords:

Freezing tolerance;
Infectivity;
Persistence study;
Trichinella;
Wild boars

*Correspondence

Email:
fernandoaf@fvvet.uba.ar

Abstract

Background: The aim of this study was to investigate the survival of *Trichinella spiralis* and *T. pseudospiralis* in decaying wild boar tissue and assess their freezing tolerance in experimentally infected animals.

Methods: The present study was conducted in Buenos Aires City, Argentina during the 2018-2019 period. Two wild boars were used, one infected with 20,000 muscle larvae (ML) of *T. spiralis* and the other with *T. pseudospiralis*. Both animals were euthanized 19 weeks post-infection. Limbs from each boar were placed over soil in plastic containers to assess ML survival in decaying tissue, under natural temperature and humidity, shielded from rain. Weekly samples were taken for artificial digestion, and the ML were inoculated into mice to determine their reproductive capacity index (RCI). Additionally, to evaluate the freezing tolerance of the ML, muscle samples were stored at -18°C. Six samples were taken and digested after 2, 4, 7, 9, 11, and 14 days, with subsequent inoculation into mice to assess RCI.

Results: *T. spiralis* remained infective in decaying wild boar tissue for 11 weeks, while *T. pseudospiralis* remained infective for only 4 weeks. The freezing tolerance assay showed that *T. spiralis* ML remain infective for 9 days. However, *T. pseudospiralis* ML remain infective for only 2 days at -18°C.

Conclusion: The findings highlight the survival strategies of *T. spiralis* and *T. pseudospiralis* in different environmental conditions, which may have implications for understanding their transmission dynamics in wild animals.



Introduction

A wide range of animals are involved in the maintenance of *Trichinella* spp. infection in wildlife (1). The probability of muscle larvae (ML) being ingested by carnivores or scavengers increases in direct proportion to the ML ability to remain viable in decaying tissue (2). Environmental conditions, such as humidity and temperature, play a key role in the spread of *Trichinella* spp., as they directly affect the survival of the larvae in host carcasses (3,4).

The ability of ML to survive in tissues exposed to low or high temperatures may be influenced by the presence or absence of a collagen capsule surrounding the ML. The collagen capsule may provide some protection and therefore increase the survival time of these larvae in decaying tissues (5 - 7). It may also confer some freeze tolerance (8). Studies have shown that the tolerance temperature of the ML is influenced by both the host species and the *Trichinella* species, among other factors (9 - 12).

Wild boars (*Sus scrofa*) are known to be a source of *Trichinella* spp. infection for humans (13 - 16). Consumption of infected meat from these animals has been found to be one of the main causes of human trichinellosis (3,14, 17 - 22).

Trichinella spp. have diverse freeze tolerances that vary depending on each species. *T. nativa* has survived in arctic fox tissues (*Alopex lagopus*) at -18 °C for up to 4 years (20), *Trichinella T6* has survived in grizzly bear tissues (*Ursus arctos*) at -6.5 to -20 °C for up to 34 months (9), and *T. britovi* has survived in laboratory rat muscles at -5 °C for up to 1 week (12). This tolerance also varies depending on the host species involved (23). Therefore, as *Trichinella* species have diverse freeze tolerances, relying solely on freezing meat as a preventive method could pose risks to food safety.

The purpose of this study was to determine whether *T. spiralis* and *T. pseudospiralis* can sur-

vive in decaying wild boar tissue and to assess the freezing tolerance of these *Trichinella* species in experimentally infected animals.

Materials and Methods

Animals

Two 60-day-old castrated wild boars were used in this study. They underwent a 30-day adaptation period to the new environment where the study was to be conducted. Clinical and hematological parameters were evaluated to assess their health condition (24). The present study was conducted in Buenos Aires City during the 2018-2019 period. All animals were housed in an indoor pigpen with plastic slatted floors at the experimental facilities of the Faculty of Veterinary Sciences, University of Buenos Aires, Argentina.

Ethics approval

The study was approved by the Institutional Committee for Use and Care of Laboratory Animals of the Faculty of Veterinary Sciences, University of Buenos Aires (CICUAL), permit number 2015/16.

One animal was orally infected with 20,000 *T. spiralis* L1 larvae (ISS1097), while the other was infected with 20,000 *T. pseudospiralis* L1 (21). These *Trichinella* species were maintained by serial passages using CF1 mice, and ML were recovered by artificial digestion (24). After 19 weeks post-infection (p.i.), the wild boars were euthanized (26). To determine the muscle larval burden in each animal, 100 g of upper forelimb and 100 g of upper hindlimb were artificially digested (26). These tissue amounts were taken from different parts of both. The recovered larvae were then inoculated into three mice to assess their infective capacity (day 0).

Persistence

One forelimb and one hindlimb were severed from each wild boar and placed over a

layer of soil in plastic containers. All containers were covered with metallic mosquito net screens, protected them from rain, exposed to natural environmental temperature and humidity conditions. The study was carried out during the Argentine autumn and winter seasons for 16 weeks. The environmental temperatures and relative humidity ranged from 9.0 to 16.6 °C and 33.75 to 99.25%, respectively. This data was obtained from the National Weather Service in Argentina (27).

Decaying muscle from limbs infected with *T. spiralis* and *T. pseudospiralis* in wild boars was analyzed weekly for 16 weeks. A 50-gram sample from each limb was taken weekly and artificially digested (26). These samples were taken from different points of both limbs due to uneven ML's distribution. Then, three BALB/c mice were orally inoculated with 300 larvae in 15 µl. After 6 weeks p.i., the mice were sacrificed, and the reproductive capacity index (RCI) was assessed to determine the infectivity of the recovered larvae (28). The RCI was calculated as the number of larvae recovered after digestion divided by the number of larvae orally inoculated.

The data about temperature and humidity were obtained from the National Weather Service in Argentina (27).

Freezing tolerance

To study the freezing tolerance of the ML of *T. spiralis* and *T. pseudospiralis*, one hindlimb and one forelimb from each wild boar were used. From these muscle groups, six 2 cm thick, 200 g samples were taken and placed in individual plastic bags. All samples were stored at -18 °C for 14 days. At each time point (2, 4, 7, 9, 11, and 14 days), a sample of each muscle group was thawed at 4 °C in the refrigerator and then artificially digested. The recovered larvae were counted, and four BALB/C mice were orally inoculated with 300 larvae. After 6 weeks pi., the mice were sacrificed, and the RCI was assessed. Both meat

samples at each point of analysis were digested in combination.

Statistical analysis

Mean RCI values for *T. spiralis* and *T. pseudospiralis* were assessed in two experiments: persistence and freezing, at different time points. Prior to ANOVA, normality of the data was confirmed via Shapiro-Wilk test. ANOVA was then employed to compare mean RCI values between species. The Tukey test compared means at significant time points ($\alpha = 0.05$). All statistical analyses were performed using In-foStat™ software.

Results

Larvae were recovered up to week 16 and week 10 from *T. spiralis* and *T. pseudospiralis*, respectively, from decaying tissues. In the freezing tolerance experiment, larvae from both *Trichinella* species were recovered after 14 days at -18 °C.

The highest mean RCI value found in the persistence study was 89.50 for *T. spiralis* in tissues decaying for 0 weeks, while for *T. pseudospiralis*, it was 75.14 in tissues decaying for 0 weeks.

The infective larvae recovered from the decaying tissues remained viable until week 11 for *T. spiralis* and week 4 for *T. pseudospiralis*, as shown in Table 1. After these periods, the RCI tested negative for both *Trichinella* species. The last positive RCI mean value obtained for decaying tissues infected with *T. spiralis* was 0.0056, and for decaying tissues infected with *T. pseudospiralis*, it was 15.48.

Although *T. spiralis* larvae were recovered from decaying wild boar tissues for 16 weeks, from the 11th week on, all larvae showed significant cuticle damage, absence of internal content, and no motility after being maintained for 1 hour at 37 °C. The RCI values from that point on were consistently negative.

Table 1: Mean RCI of the mice infected with *T. spiralis* and *T. pseudospiralis* ML recovered from decaying muscle tissue

Weeks	RCI		F ^b	P value
	<i>T. spiralis</i>	<i>T. pseudospiralis</i>		
0	Mean +/- SE ^a 89.51 +/- 5.73 A*	Mean +/- SE 75.14 +/- 5.73 A	3.14	0.151
1	64.93 +/- 5.98 A	69.47 +/- 5.98 A	0.29	0.6205
2	50.46 +/- 5.17 A	57.07 +/- 5.17 A	0.82	0.4176
3	48.34 +/- 3.52 A	50.96 +/- 3.52 A	0.28	0.6262
4	47.98 +/- 3.35 A	15.49 +/- 3.35 B	46.99	0.0024
5	36.98 +/- 2.02 A	0 +/- 2.02 B	167.64	0.0002
6	34.70 +/- 3.69 A	0 +/- 3.69 B	44.25	0.0027
7	23.01 +/- 4.44 A	0 +/- 4.44 B	13.42	0.0215
8	5.46 +/- 1.9 A	0 +/- 1.9 A	4.12	0.1123
9	1.58 +/- 0.39 A	0 +/- 0.39 B	8.22	0.0456
10	0.0036 +/- 0.0026 A	0 +/- 0.0026 A	1	0.3739
11	0.0056 +/- 0.0026 A	0 +/- 0.0026 A	1	0.3739

^aStandard Error

^bF critical values

*Means with a common letter are not significantly different (*P*-value > 0.05).

Regarding the freezing tolerance study, the mice infected with *T. spiralis* ML obtained from frozen muscles (-18°C) remained infectious for 9 days (Table 2), while the mice infected with frozen *T. pseudospiralis* ML remained infectious for up to 2 days at -18°C (Table 2). *T. pseudospiralis* larvae remained motile after digestion in samples frozen for up to 2 days, but became immotile thereafter. Both *Trichinella* species recovered from samples fro-

zen for 11 and 14 days showed severe cuticle damage.

The statistical analysis showed significant differences between the RCI mean values for *T. spiralis* and *T. pseudospiralis* for the persistence experiment in weeks 4, 5, 6, 7, and 9 (Table 1). For the freezing experiment, the analysis showed significant differences between the RCI mean values of *T. spiralis* and *T. pseudospiralis* for days 2, 4, and 7 (Table 2).

Table 2: Mean RCI obtained from the mice infected with *T. spiralis* and *T. pseudospiralis* ML recovered from frozen muscle tissue, after different times under freezing conditions

Days	RCI		F ^b	P-value
	<i>T. spiralis</i>	<i>T. pseudospiralis</i>		
0	Mean +/- SE ^a 89.51 +/- 5.73 A*	Mean +/- SE 75.14 +/- 5.73 A	3.14	0.151
2	52.84 +/- 6.53 A	17.45 +/- 6.53 B	14.68	0.0186
4	30.42 +/- 3.92 A	0 +/- 3.92 B	30.13	0.0054
7	12.65 +/- 2.82 A	0 +/- 2.82 B	10.1	0.0336
9	1.93 +/- 0.97 A	0 +/- 0.97 A	1.96	0.2341

^aStandard Error

^bF critical values

*Means with a common letter are not significantly different (*P*-value > 0.05)

Discussion

In the present study, we observed that the ability of ML to survive in decaying and frozen muscle tissue varied according to the *Trichinella* species. Specifically, *T. spiralis* exhibited a longer survival time in both decaying and frozen meat than *T. pseudospiralis*.

The age of the ML has been suggested as a factor that can affect the ML ability to survive in decaying tissue, as previously reported by several authors (5-7). In a study (7), mice infected with *T. spiralis* for 5 weeks had ML surviving in decaying tissue for 1 week, while those infected for 37 weeks had ML surviving for up to 3 weeks under the same conditions. Prolonged infection time may result in the collagen capsule becoming thicker and more fully developed, which could favor encapsulated species like *T. spiralis*. In this study, wild boars were infected for over 19 weeks, and the *T. spiralis* ML remained infective for almost 11 weeks in decaying wild boar tissues, while *T. pseudospiralis* ML only remained infective for 4 weeks in decaying tissues. This finding supports the idea that the age of the ML capsule may be beneficial for its survivability in decaying tissues. In a study conducted in guinea pigs (*Cavia porcellus*) decomposing tissues infected with *T. patagoniensis* (33), the ML could survive for several weeks. In addition, several authors suggest that high temperatures and droughts reduce the survival capacity of these encysted larvae (34, 35).

In our experiment, the ML of *T. pseudospiralis* in decaying tissue were infective for 4 weeks. The shorter survival time observed in *T. pseudospiralis* compared to *T. spiralis* may be attributed to the fact that *T. pseudospiralis* is a non-encapsulated species. Several authors have suggested that the presence of a collagen capsule surrounding the ML benefits its survival in carcasses (5 - 6, 29). Therefore, the increased susceptibility of *T. pseudospiralis* to putrefaction byproducts could be a conse-

quence of the absence of the collagen capsule, which provides greater protection (30 - 31).

Studies conducted with *T. papuae*, a non-encapsulated species, have shown varying results in terms of its survival in carcasses. For instance, its ML could survive for up to 9 days in domestic pig carcasses under the environmental conditions of Papua New Guinea (2). Similarly, it was demonstrated that *T. papuae* remained infective for more than 9 days at a temperature above 20 °C in red fox (*Vulpes vulpes*) carcasses (32).

According to Pozio (36), various factors affect the ability of *Trichinella* ML to survive freezing temperatures, including the host in which they are encysted, the ML age (12, 37), the temperature and time exposed (37, 38). Some *Trichinella* species have been found to survive longer in carnivorous muscles than in pigs or laboratory rodent muscles (37, 39 - 41). Many authors suggest that differential muscle composition, such as the type and amount of lipids present in the tissue, may provide some protection against freezing temperatures in wild animals since they face different environmental conditions than domestic animals (42). Additionally, the ML ability to tolerate freezing temperatures may also be related to the levels of endogenous carbohydrates present in muscle tissue and on larvae glycogen concentrations (43, 44). The temperature range is also a key factor in the ML's ability to survive in frozen tissues. The optimal temperature range of survival is 0 to -20 °C, specifically the temperature "below the snow" (9, 34, 35, 45). Finally, the ML survival capacity in freezing environments may be directly linked to the genetic differences between *Trichinella* species (36).

In our experiment, we observed that *T. pseudospiralis* larvae had a low tolerance for freezing, as they only survived for 2 days at -18 °C. (31). Similarly, in a previous study on shredded rat tissues it was reported that *T. pseudospi-*

ralis could survive for less than one week at a freezing temperature of -18°C (12). In contrast, it was demonstrated (46) that *T. pseudospiralis* ML in sheep muscle tissue remained infective for up to 4 weeks at -18°C .

Differences in the persistence and infectivity of *T. pseudospiralis* populations have been observed in various animal hosts (10,40,41). *T. pseudospiralis* has been found in 18 different mammal species and 12 avian species, as reported by Pozio (36, 48). This wide host range could explain how the species has been able to expand and spread worldwide despite its low tolerance to different environmental conditions (36, 50).

T. spiralis is the *Trichinella* species that is best adapted to domestic and wild swine hosts (49, 51, 52). Kapel et al. (44) studied nine *Trichinella* species and found that none survived over one week at -18°C in pig or wild boar meat. Another study in wild boars showed a survival rate for 56 hours at -21°C (8). However, our study found that *T. spiralis* could survive for 9 days at -18°C .

In the present study, decaying limbs were exposed to autumn and winter environmental conditions, which allowed the larvae of *T. spiralis* to survive for 11 weeks and *T. pseudospiralis* for 4 weeks. However, in frozen tissue, *T. spiralis* only survived for 9 days and *T. pseudospiralis* for 2 days. Therefore, the ability to survive in decaying and frozen tissue indicates that these two *Trichinella* species can survive in carcasses and be spread by scavenger animals. These findings are crucial for understanding the epidemiology of *Trichinella* spp., especially given the importance of wild boars as an exotic species widely distributed in Argentina and America.

Conclusion

The findings highlight the survival strategies of *T. spiralis* and *T. pseudospiralis* in different environmental conditions, which may have

implications for understanding their transmission dynamics in wild animals.

Acknowledgements

This work was supported by UBACyT 20020170100530BA, 20020190200396BA, MINCyT, FONCyT PICT-2015-2350 and PICT-2018-01203.

Conflict of Interest

The authors declare that there is no conflict of interests.

References

1. Santrac V, Nedic DN, Maric J, et al. The first report of *Trichinella pseudospiralis* presence in domestic swine and *T. britovi* in wild boar in Bosnia and Herzegovina. *Acta Parasitol.* 2015; 60(3):471-475.
2. Owen IL, Reid SA. Survival of *Trichinella papuae* muscle larvae in a pig carcass maintained under simulated natural conditions in Papua New Guinea. *J Helminthol.* 2007; 81:429-432.
3. Pozio E, Darwin Murrell K. Systematics and Epidemiology of *Trichinella*. *Adv Parasitol.* 2006; 63:367-439.
4. Bilska-Zajac E, Rózycki M, Grądziel-Krukowska K, et al. Diversity of *Trichinella species* in relation to the host species and geographical location. *Vet Parasitol.* 2020; 279:109052.
5. Campbell WC. *Trichinella* and Trichinellosis. Plenum Press, New York, 1983. p. 581.
6. Stewart GL. Pathophysiology of the Muscle Phase, in: Campbell, W.C. (Ed.), *Trichinella* and Trichinosis. Springer US, Boston, MA. 1983. p. 241-264.
7. Von Köller J, Kapel CMO, Enemark HL, Hindsbo O. Infectivity of *Trichinella* spp. recovered from decaying mouse and fox muscle tissue. *Parasite.* 2001; 8(2 Suppl):S209-12.
8. Lacour SA, Heckmann A, Macé P, et al. Freeze-tolerance of *Trichinella* muscle larvae in experimentally infected wild boars. *Vet Parasitol.* 2013; 194:175-178.

9. Worley DE, Seese FM, Espinosa RH, Sterner MC. Survival of sylvatic *Trichinella spiralis* isolates in frozen tissue and processed meat products. J Am Vet Med Assoc. 1986; 189:1047–1049.
10. Dick TA, Belosevic M. Observations on a *Trichinella spiralis* Isolate from a Polar Bear. J Parasitol. 1978; 64:1143–1145.
11. Pozio E, la Rosa G, Rossi P, Murrell KD. Biological Characterization of *Trichinella* Isolates from Various Host Species and Geographical Regions. J Parasitol. 1992; 78:647–653.
12. Malakauskas A, Kapel CMO. Tolerance to Low Temperatures of Domestic and Sylvatic *Trichinella* Spp. in Rat Muscle Tissue. J Parasitol. 2003; 89:744–748.
13. Burke R, Masuoka P, Murrell KD. Swine *Trichinella* infection and geographic information system tools. Emerg Infect Dis. 2008; 14:1109–1111.
14. Murrell KD, Pozio E. Worldwide occurrence and impact of human trichinellosis, 1986–2009. Emerg Infect Dis. 2011; 17:2194–2202.
15. Franssen F, Swart A, vander Giessen J, Havelaar A, Takumi K. Parasite to patient: A quantitative risk model for *Trichinella* spp. in pork and wild boar meat. Int J Food Microbiol. 2017;241:262–275.
16. Gondek M, Grzelak S, Pyz-Łukasik R, Knysz P, Ziomek M, Bień-Kalinowska J. Insight into *Trichinella britovi* infection in pigs: Effect of various infectious doses on larvae density and spatial larvae distribution in carcasses and comparison of the detection of Anti-*T. britovi* IgG of Three Different Commercial ELISA Tests and Immunoblot Assay. Pathogens. 2022; 11(7):735.
17. Dupouy-Camet J. Trichinellosis: A worldwide zoonosis. Vet Parasitol. 2000; 93:191–200.
18. Ribicich M, Gamble HR, Bolpe J, et al. *Trichinella* infection in wild animals from endemic regions of Argentina. Parasitol Res. 2010; 107:377–380.
19. Fichi G, Stefanelli S, Pagani A, et al. Trichinellosis Outbreak Caused by Meat from a Wild Boar Hunted in an Italian Region Considered to be at Negligible Risk for *Trichinella*. Zoonoses Public Health. 2015; 62(4):285–291.
20. Kapel CMO, Pozio E, Sacchi L, Prestrud P. Freeze Tolerance, Morphology, and RAPD-PCR Identification of *Trichinella nativa* in Naturally Infected Arctic Foxes. J Parasitol. 1999; 85: 144–147.
21. Rostami A, Gamble HR, Dupouy-Camet J, Khazan H, Bruschi F. Meat sources of infection for outbreaks of human trichinellosis. Food Microbiol. 2017; 64:65–71.
22. Faber M, Schink S, Mayer-Scholl A, et al. Outbreak of trichinellosis due to wild boar meat and evaluation of the effectiveness of post exposure prophylaxis, Germany, 2013. Clin Infect Dis. 2015; 60(12):e98–e104.
23. Hill DE, Forbes L, Gajadhar AA, Gamble HR. Viability and infectivity of *Trichinella spiralis* muscle larvae in frozen horse tissue. Vet Parasitol. 2007; 146:102–106.
24. Bessi C, Ercole ME, Fariña FA, et al, Pasqualetti MI. Study of *Trichinella patagoniensis* in wild boars. Vet Parasitol. 2021; 297:109166.
25. Krivokapich SJ, Gonzalez-Prous CL, Gatti GM, Saldía L. First finding of *Trichinella pseudospiralis* in the Neotropical region. Vet Parasitol. 2015; 208:268–271.
26. Gamble HR, Bessonov AS, Cuperlovic K, et al. International Commission on Trichinellosis: Recommendations on methods for the control of *Trichinella* in domestic and wild animals intended for human consumption. Vet Parasitol. 2000; 93:393–408.
27. Anonymous. Servicio Meteorológico Nacional. Observatorio Buenos Aires. 2022. <https://www.estadisticaciudad.gob.ar/eyc/?p=27702>. Accessed 11 January 2022.
28. Kapel CMO, Gamble HR. Infectivity, persistence, and antibody response to domestic and sylvatic *Trichinella* spp. in experimentally infected pigs. Int J Parasitol. 2000; 30:215–221.
29. Riva E, Steffan P, Guzmán M, Fiel C. Persistence of *Trichinella spiralis* muscle larvae in natural decaying mice. Parasitol Res. 2012; 111:249–255.
30. Stewart GL, Kennedy R, Larsen E. Infectivity of *Trichinella pseudospiralis* Isolated from Carrion, Source: J Parasitol. 1990; 76(5):750–751.
31. Kapel CMO. Host diversity and biological characteristics of the *Trichinella* genotypes and their effect on transmission. Vet Parasitol. 2000; 93:263–278.
32. Webster P, Malakauskas A, Kapel CMO. Infectivity of *Trichinella papuae* for

- experimentally infected red foxes (*Vulpes vulpes*). Vet Parasitol. 2002; 105:215–218.
33. Fariña F, Pasqualetti M, Ilgová J, Cardillo N, et al. Evaluation of the infectivity and the persistence of *Trichinella patagoniensis* in muscle tissue of decomposing guinea pig (*Cavia porcellus*). Parasitol Res. 2017; 116:371-375.
 34. Pozio E. The impact of globalization and climate change on *Trichinella* spp. epidemiology. Food Waterborne Parasitol. 2022; 27:e00154.
 35. Keppel G, Wardell-Johnson GW. Refugia: keys to climate change management. Glob Chang Biology. 2012; 18(8):2389-2391.
 36. Pozio E. Adaptation of *Trichinella* spp. for survival in cold climates. Food Waterborne Parasitol. 2016; 4:4–12.
 37. Pozio E, La Rosa G, Amati M. Factors influencing the resistance of *Trichinella* muscle larvae to freezing. In: Trichinellosis. Proceedings of the 8th International Conference on Trichinellosis. Ed. Campbell CW, Pozio E, and Bruschi F. Rome: Istituto Superiore di Sanita Press, 1994. p:173–178.
 38. Sacchi L, Corona S, Gajadhar AA, Pozio E. Ultrastructural characteristics of nurse cell-larva complex of four species of *Trichinella* in several hosts. Parasite. 2011; 8(2 Suppl):S54-8.
 39. Kapel CMO, Webster P, Malakauskas A, Hurnikova Z, Gamble HR. Freeze tolerance of *Trichinella* genotypes in muscle tissue of experimentally infected pigs, horses, wild boars, mice, cats, and foxes. In: International Conference Trichinellosis, San Diego, 2004. p. 8-12.
 40. Seymour J, Horstmann-Dehn L, Rosa C, Lopez JA. Occurrence and genotypic analysis of *Trichinella* species in Alaska marine-associated mammals of the Bering and Chukchi seas. Vet Parasitol. 2014; 200(1-2):153-164.
 41. Rudneva OV, Andreyanov ON, Sidor EA. Reprint of: The retention and concentration of glycogen in *Trichinella nativa* in the winter-spring period. Vet Parsitol. 2021; 297:109547.
 42. Dick TA. Infectivity of isolates of *Trichinella* and the ability of an arctic isolate to survive freezing temperatures in the raccoon, *Procyon lotor*, under experimental conditions. J Wildl Dis. 1983; 19:333–336.
 43. Raines KM, Stewart GL. Carbohydrate reserves and infectivity in *Trichinella spiralis* isolated from carrion. Parasitology. 1988; 96(Pt 3):533-41.
 44. Davidson RK, Handeland K, Kapel CM. High tolerance to repeated cycles of freezing and thawing in different *Trichinella nativa* isolates. Parasitol Res. 2008; 103:1005-1010
 45. Pauli JN, Zuckerberg B, Whiteman JP, Porter W. The subnivium: a deteriorating seasonal refugium. Front Ecol Environ. 2013; 11:260-267.
 46. Theodoropoulos G, Kapel CMO, Webster P, Saravanos L, Zaki J, Koutsotolis K. Infectivity, predilection sites, and freeze tolerance of *Trichinella* spp. In experimentally infected sheep. Parasitol Res. 2000; 86:401–405.
 47. Malakauskas A, Kapel CM., Webster P. Infectivity, persistence and serological response of nine *trichinella* genotypes in rats. Parasite. 2001; 8(2 Suppl):S216-22.
 48. Pozio E. The broad spectrum of *Trichinella* hosts: From cold- to warm-blooded animals. Vet Parasitol. 2005; 132:3–11.
 49. Gondek M, Knysz P, Pomorska-Mól M, Ziomek M, Bień-Kalinowska J. Acute phase protein pattern and antibody response in pigs experimentally infected with a moderate dose of *Trichinella spiralis*, *T. britovi*, and *T. pseudospiralis*. Vet Parasitol. 2020; 288:109277.
 50. Pozio, E. *Trichinella pseudospiralis* an elusive nematode. Vet Parasitol. 2016; 231:97–101.
 51. Gottstein B, Pozio E, Nöckler K. Epidemiology, diagnosis, treatment, and control of trichinellosis. Clin Microbiol Rev. 2009; 22(1):127-145.
 52. Pozio E, Zarlenga DS. Recent advances on the taxonomy, systematics and epidemiology of *Trichinella*. Int J Parasitol. 2005; 35:1191–1204.